



OPEN ACCESS

EDITED BY

Claudia Spadavecchia,
University of Bern, Switzerland

REVIEWED BY

Stelio Pacca Loureiro Luna,
Sao Paulo State University, Brazil
Mathieu De Preux,
University of Bern, Switzerland

*CORRESPONDENCE

Ulrike Auer
✉ ulrike.Auer@vetmeduni.ac.at
Florien Jenner
✉ florien.Jenner@vetmeduni.ac.at

[†]These authors have contributed equally to this work

RECEIVED 11 September 2023

ACCEPTED 08 December 2023

PUBLISHED 19 January 2024

CITATION

Auer U, Kelemen Z, Vogl C, von Ritgen S, Haddad R, Torres Borda L, Gabmaier C, Breteler J and Jenner F (2024) Development, refinement, and validation of an equine musculoskeletal pain scale.
Front. Pain Res. 4:1292299.
doi: 10.3389/fpain.2023.1292299

COPYRIGHT

© 2024 Auer, Kelemen, Vogl, von Ritgen, Haddad, Torres Borda, Gabmaier, Breteler and Jenner. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Development, refinement, and validation of an equine musculoskeletal pain scale

Ulrike Auer^{1*†}, Zsolia Kelemen², Claus Vogl³,
Stephanie von Ritgen¹, Rabea Haddad², Laura Torres Borda²,
Christopher Gabmaier¹, John Breteler² and Florian Jenner^{2*†}

¹Anaesthesiology and Perioperative Intensive Care Medicine Unit, Department of Companion Animals and Horses, University of Veterinary Medicine Vienna, Vienna, Austria, ²Equine Surgery Unit, Department of Companion Animals and Horses, University Equine Hospital, University of Veterinary Medicine Vienna, Vienna, Austria, ³Department of Biomedical Sciences, Institute of Animal Breeding and Genetics, University of Veterinary Medicine Vienna, Vienna, Austria

Musculoskeletal disease is a common cause of chronic pain that is often overlooked and inadequately treated, impacting the quality of life of humans and horses alike. Lameness due to musculoskeletal pain is prevalent in horses, but the perception of pain by owners is low compared with veterinary diagnosis. Therefore, this study aims to establish and validate a pain scale for chronic equine orthopaedic pain that is user-friendly for horse owners and veterinarians to facilitate the identification and monitoring of pain in horses. The newly developed musculoskeletal pain scale (MPS) was applied to 154 horses (mean age 20 ± 6.4 years SD) housed at an equine sanctuary, of which 128 (83%) suffered from chronic orthopaedic disease. To complete the MPS, the horses were observed and videotaped from a distance while at rest in their box or enclosure. In addition, they received a complete clinical and orthopaedic exam. The need for veterinary intervention to address pain (assessed and executed by the sanctuary independent from this study) was used as a longitudinal health outcome to determine the MPS's predictive validity. To determine the interrater agreement, the MPS was scored for a randomly selected subset of 30 horses by six additional blinded raters, three equine veterinary practitioners, and three experienced equestrians. An iterative process was used to refine the tool based on improvements in the MPS's correlation with lameness evaluated at the walk and trot, predictive validity for longitudinal health outcomes, and interrater agreement. The intraclass correlation improved from 0.77 of the original MPS to 0.88 of the refined version (95% confidence interval: 0.8–0.94). The refined MPS correlated significantly with lameness at the walk ($r = 0.44$, $p = 0.001$) and trot ($r = 0.5$, $p < 0.0001$). The refined MPS significantly differed between horses that needed veterinary intervention (mean MPS = 8.6) and those that did not (mean MPS = 5.0, $p = 0.0007$). In summary, the MPS showed good interrater repeatability between expert and lay scorers, significant correlation with lameness at the walk and trot, and good predictive validity for longitudinal health outcomes, confirming its ability to identify horses with orthopaedic health problems.

KEYWORDS

chronic pain, pain scale, musculoskeletal pain, discomfort, horse, equine

1 Introduction

Musculoskeletal disease is the leading cause of chronic pain in horses and humans alike (1–8). In equine veterinary practice, lameness due to musculoskeletal pain ranks as the most prevalent diagnosis (1–4, 9–12). Already, in 4- to 5-year-old riding horses, 24% demonstrated moderate to severe orthopaedic clinical findings (12), emphasizing the widespread nature of the problem. The prevalence further increases in older horses, with 51% of horses above 15 years and 77% of horses aged 30 years and older exhibiting lameness, which is strongly associated with pain experienced at rest (3–5, 11, 13).

Despite their high prevalence, musculoskeletal diseases are frequently overlooked as a source of suffering and, as a result, receive inadequate treatment (2–4, 14–16). Indeed, owners reported lameness in only 16% of horses compared with the 77% diagnosed by veterinarians in the same cohort (11). Similarly, in two other groups of horses in training that were perceived to be sound by their owners, 72.5% and 74% showed movement asymmetry during objective lameness evaluation (17, 18). The owners' low perception of musculoskeletal pain compared with the expert diagnoses is concerning from both a veterinary and welfare perspective. It further compounds the undertreatment of pain also observed in older humans that is associated with the erroneous but widespread societal belief that pain is a natural part of ageing and inevitable in later life (15, 19).

Due to the subjective nature of pain, *gold standard* pain assessment tools in human medicine rely on self-reporting, as direct measurement of individual experiences is not feasible (20, 21). For patients unable to communicate in ways easily understood by their caregivers, such as non-verbal human patients and animals, pain assessment depends on physiological and behavioural indicators (21–27). However, physiologic indicators, including changes in heart and respiratory rate, lack the sensitivity and specificity needed for reliable pain detection and discrimination from other sources of distress (28, 29). Although these indicators are commonly used to indicate the presence of pain, little empirical evidence exists to support this practice, as the correlation of vital sign changes with self-reports is weak, and the absence of changes in vital signs does not necessarily mean the absence of pain (28, 29).

By contrast, research has shown a strong correlation between pain behaviours and patients' verbal pain reports, though external observers tend to underestimate pain intensity (30, 31). Consequently, non-verbal pain behaviours, such as facial expressions, lameness, and guarding, have become integral to pain assessments (22, 24, 27–42). Especially facial expressions, which have been demonstrated to encode both the sensory and affective components of pain, are commonly used to recognize and quantify pain in human and veterinary patients who are unable to verbalize (13, 26, 27, 37–50). Postural and gait adaptations that reduce the load on painful tissue to prevent or alleviate pain and protect from further injury (51) are also strongly associated with orthopaedic pain in humans and horses alike (35, 39, 40). However, despite evidence that guarding and posture may be more indicative of musculoskeletal pain than

facial expressions (35, 39, 40), body cues are not routinely included in pain assessment.

As some behavioural changes associated with chronic pain may develop gradually and be subtle, making them most easily detected by someone familiar with the animal and its behaviour before and after the onset of pain (26, 52), the inclusion of caretaker assessments can add important cues to facilitate identification of equine pain. Regular pain assessment by caretakers is also essential to optimize treatment, as chronic musculoskeletal conditions typically require prolonged and often life-long palliative treatment and therapy adjustments to address acute flares and fluctuations in pain intensity while minimizing side effects. Hence, there is a clear need for a pain assessment tool that horse owners and veterinarians alike can use to facilitate the identification of pain, communication between veterinarians and clients, and the evaluation of the effectiveness of pain management interventions. This pain scoring system should be based on objective measures that are sensitive and specific to pain and minimize the potential for observer bias and misinterpretation (20, 23). Keeping in mind horses' instinctive tendency to exhibit little indication of pain in the presence of potential predators, such as humans, and to reduce or relieve pain behaviour even during caretaker visits (46), the pain assessment tool should also be applicable remotely using video surveillance or recordings.

Therefore, this study aims to establish, refine, and validate an orthopaedic pain scale that is easy and fast to use by horse caretakers and veterinarians alike, and can also be used to score pain on videos to minimize observer interference with pain behaviour. Based on recent scientific evidence, the newly developed equine musculoskeletal pain scale (MPS) incorporates components of the equine pain face (27, 38), posture, head-neck position, weight-bearing, and weight shifting to assess orthopaedic pain in horses (33, 36, 39, 40, 44, 47, 53).

2 Materials and methods

This prospective, observational cohort study was designed to refine and validate the newly developed MPS, a tool based on components of the equine pain face (38), recent scientific advances demonstrating the importance of posture, weight-bearing, and head position for chronic pain behaviour (5, 13, 33, 36, 40, 44–47, 53–57), and clinical observations in patients suffering from chronic orthopaedic pain. During scale development and refinement, a panel of six experts (three equine veterinarians and three experienced equestrians) assessed content validity and comprehensibility through iterative evaluations of item relevance, comprehensiveness, and clarity (see Sections 2.3 and 2.5) (58–63). The size of the expert panel was based on previous studies establishing a minimum of four to five experts to be adequate for content validation (63, 64). In addition, item relevance was assessed by calculating the correlation between each item and the total MPS and by an item–total correlation (see Sections 2.3 and 2.7). To evaluate the influence of different sources of variability on the MPSs, reliability was determined by calculating the interrater variability in relation to the horses'

variability and the total variability in a mixed model (see Sections 2.3, 2.5, and 2.7) (58–63). The MPS's construct validity was assessed by calculating the correlation of the MPS with subjective lameness scores at the walk and trot and the objective lameness data (see Section 2.4) (58–63). The scale's criterion validity to predict longitudinal health outcomes was evaluated by comparing the MPS score of horses that required veterinary intervention in the subsequent months with those that did not need medical treatment (see Section 2.4) (58–63). Lastly, the scale's responsiveness was assessed by comparing the MPSs of horses that received analgesia before and after treatment (58–63).

2.1 Horses

A total of 154 horses living at an equine sanctuary were included in this study. The horses were maintained in their familiar environment and husbandry conditions, and neither the horses' housing, turn-out, or feeding regime nor any other management factors or veterinary treatments were affected by the study or changed for study purposes.

Before inclusion in the study, all horses underwent an in-depth physical exam. The horses suffering from non-orthopaedic causes of pain or cardiovascular (e.g., ventricular tachycardia) or gastrointestinal (e.g., delayed gastric emptying) disease were excluded from the study.

2.2 Horse examination and assessment parameters

All the horses were examined by the same veterinarian and received a complete clinical and orthopaedic exam in addition to an MPS. To complete the MPS, the horses were discreetly observed and videotaped from a suitable distance (5–10 m), displaying no awareness or curiosity towards the observer, while at rest in their box or enclosure (paddock or pasture). The MPS uses an ordinal scale to measure demeanour (13, 25, 26, 40, 44–47, 49, 65–67), pain face (22, 27, 38, 68, 69), weight shifting, weight-bearing, head-neck posture, limb posture (25, 33, 36, 44, 47, 53, 56, 57), and lameness that is evident while observing the horse from a distance in its enclosure (Supplementary Figure S1). Examinations involving direct interaction with the horses were conducted only after all initial distant observations were finished. This approach was taken to minimize any potential influence on the horses' behaviour and any biasing of the MPS results by the examination process.

The orthopaedic exam included a subjective lameness evaluation grading the lameness at the walk and trot separately on a scale from 0 (sound) to 5 (non-weight-bearing) (70). The horses that were unable to trot because of severe lameness were assigned a score of 5 for the lameness at the trot. Horses with a lameness score >2 at the walk and ≥ 3 at the trot were considered moderately to severely lame.

In addition, the lameness was assessed objectively in 110 horses (71.4%) using a commercially available multi-sensor inertial gait analysis system (Lameness Locator[®], Equinosis, USA) that has

been validated to detect and quantify equine lameness (71–76). The horses were considered lame with a Q-score, a metric quantifying movement asymmetry amplitude, >8.5 mm and moderately to severely lame with a Q-score >30 mm (74–80).

2.3 Musculoskeletal pain scale—descriptive statistics

To characterize the MPS and its items, a correlation matrix was calculated among pairs of scores of items and the total using the non-parametric Spearman and the parametric Pearson correlations. In addition, a principal component analysis was calculated to further characterize the relationships among items. Both methods allow evaluation of which variables contribute independently or jointly to the total MPS.

2.4 Musculoskeletal pain scale—validity and predictive performance

The primary measures for assessing the validity of the MPS were the subjective and objective lameness scores. The correlations between the MPS with subjective lameness at the walk and trot and the objective lameness data were calculated using moderate to severe lameness as an indicator of pain.

To assess the MPS's predictive performance for longitudinal health outcomes, the MPSs of horses that required veterinary intervention [analgesia (firocoxib or phenylbutazone) or euthanasia for pain that was unresponsive to treatment] in the months following the exam were compared retrospectively with those that did not need medical treatment. The need for veterinary intervention was determined by the nursing and veterinary staff of the sanctuary based on their independent assessment of the horses' health and pain status, thereby providing an outcome variable independent from the study.

2.5 Musculoskeletal pain scale—interrater agreement and refinement

To ensure the inclusion of horses representing the entire spectrum of pain grades in the interrater agreement analysis, the horses were considered pain-free if their MPS was ≤ 3 ($n = 64$), mildly painful if their MPS was between 4 and ≤ 8 ($n = 69$), and moderately to severely painful if their MPS was ≥ 9 ($n = 20$). A subset of 30 horses, 10 of each pain group, was randomly selected using the GraphPad[®] random selection tool (<https://www.graphpad.com/quickcalcs/randomselect2/>). To determine the interrater agreement, six additional raters, three equine veterinary practitioners and three experienced equestrians, completed the MPSs for these 30 horses. The six additional raters were blinded to the horses' medical history, pain group, and exam results and completed the MPS based solely on anonymized videos obtained during the exam. Interrater reliability was assessed using intraclass correlation (ICC) analysis.

Based on the interrater agreement and their feedback regarding the clarity of the item descriptions and the scoring process, the MPS was refined to optimize the discriminative power of the items to ensure unequivocal definitions of each item to limit the potential for misinterpretation and to shorten the time required to complete the MPS to enhance its clinical and research utility. An iterative process was used in tool refinement, considering improvements achieved (content, construct, and criterion validity, comprehensiveness, comprehensibility, reliability, interrater agreement) compared with the original tool when replacing existing items or adding items. Item redundancy was investigated using correlation and principal component analysis. The refined MPS measures seven items on an ordinal scale and can accumulate a maximum score of 26 points, 2 for demeanour, 4 for a pain face, 2 for head-neck posture, 4 for weight shifting, 6 for limb posture, 4 for weight-bearing, and 4 for lameness that is evident while observing the horse from a distance in its enclosure (English version: [Figure 1](#), German version: [Supplementary Figure S2](#)). The refined tool was tested in a new randomly selected subset of 30 horses representing the three pain groups ($n = 10$ per group) to assess the interrater agreement using ICC analysis and with a mixed model with rater-ID and horse-ID as random variables (see also Section 2.7). For the ICC, interrater agreement was considered to be very good (for scores 0.81–1.0), good (0.61–0.80), moderate (0.41–0.60), reasonable (0.21–0.4), or poor (<0.2) (81). Based on the excellent interrater agreement for the MPS's lameness item established in the first validation step (ICC score 0.83), only observers one and two rated the lameness item, which had not been changed during the refinement process, as part of the last iteration of the MPS.

2.6 Refined musculoskeletal pain scale—validity and discriminative power

To assess the validity and discriminative power of the refined MPS, the refined MPS was completed for 60 video-recorded behavioural observations that had not been included in the first validation step (see Section 2.4). The construct validity was determined by correlation analysis of the MPS with the subjective lameness at the walk and the trot. The refined MPS's criterion validity to predict longitudinal health outcomes was calculated by comparing the MPSs of the horses that required veterinary intervention with those that did not, using a Mann-Whitney U test. The MPS's responsiveness, its ability to discriminate between before and after treatment, was assessed by comparing the MPSs of the horses receiving analgesia before and after intervention.

Receiver operating characteristic (ROC) analysis was utilized to evaluate the global performance of the MPS in discriminating between lame and sound horses and between horses that needed veterinary intervention and those that did not. In addition, the ROC was used to determine cut-off values that minimize misclassification errors (82–84). The optimal cut-off for discriminating between horses suffering from pain and those without any pain was identified as the value where the sum of

the sensitivity [=probability of a positive test outcome in a horse that is in pain (true-positive)] and specificity [=probability of a negative test outcome in a pain-free horse (true-negative)] was maximized. If two cut-off values yielded similar sums of sensitivity and specificity, the cut-off with the higher sensitivity was chosen to maximize the likelihood of identifying horses suffering from pain for further diagnostics and therapy if required. Since the MPS item lameness can only be assessed if the horse is moving in its stall or enclosure without being prompted, scoring this item may not always be possible, which could result in a lower maximum MPS. Therefore, we determined the cut-off value for the refined MPS, both including and excluding the lameness item.

2.7 Statistical analyses

Statistical analyses were carried out using GraphPad Prism (version 10.0.2, GraphPad Software LLC, Boston, MA, USA), NCSS 2020 Statistical Software (NCSS, LLC, Kaysville, UT, USA), and the “R” statistical programming language (R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org/>) (85). The D'Agostino–Pearson, Shapiro–Wilk, Anderson–Darling, and Kolmogorov–Smirnov tests were computed to assess whether data were normally distributed. The t -test, ANOVA, and Pearson's correlation test were used for normally distributed parameters, whereas for parameters that were not normally distributed, the Mann–Whitney U , Kruskal–Wallis, and the Spearman correlation tests were calculated. A principal component analysis was performed to describe the relationships among the MPS items. For the item–total correlation, the value of the focal item was correlated with the MPS minus the value of the focal item. Furthermore, a mixed model with horse and rater as random effects was computed to evaluate the relative contribution of each to the total variation. The p -values < 0.05 were considered statistically significant. For correlation analysis, correlation coefficients $|r| < 0.3$, $0.3 \leq |r| \leq 0.8$, and $|r| > 0.8$ were considered to indicate weak, significant, and strong correlations, respectively. For ROC analysis, the concordance statistic (c -statistic, equivalent to the area under the ROC curve) represents the probability that a randomly selected patient will have a higher test result than a randomly selected control. It is utilized as a measure of the global accuracy of a diagnostic test and is considered to indicate low, moderate, and high test accuracy at values of $0.5 < c \leq 0.7$, $0.7 < c \leq 0.9$, and $c > 0.9$, respectively (82–84).

3 Results

3.1 Horses

The 154 horses included 67 warmbloods, 25 draft, 25 Arabian, 18 Haflinger horses, and 19 horses of other breeds. The horses were 2–32 years old [mean: 20 years, SD: 6.4 years, median: 21 years, interquartile range (IQR): 16–26 years]. Of the 154 horses, 128 (83%) suffered from chronic orthopaedic disease, such as

Discomfort / Pain score®



To be completed while observing the horse (in its stall, paddock,...) from a distance!

Horse: _____ Date: _____

Demeanor (0-2 points)	Bright, alert and responsive: interested in and attentive to surroundings, ears move inquisitively, appropriate interactions with and responses to environmental stimuli e.g., ears moving in response to noises		0
	Quiet, alert and responsive: appropriate interactions with and responses to environmental stimuli		0
	Dozing/sleeping: standing quietly, head may be low, eyelids and lips may droop, response to environmental stimuli can be delayed		0
	Depressed/withdrawn: eyes open but dull and unfocused gaze, standing motionless, ears not moving, indifferent and unresponsive to environmental stimuli		2
Pain face (0-4 points)		No pain face: the muzzle, nostrils, chewing muscles and muscles above the eye are relaxed, the nostrils are elongated comma-shaped	0
		Signs of a pain face are present (e.g., dilated nostrils, tense muscles above the eyes, tense chewing muscles, ears flattened, lips pressed together) but disappear in response to the presence of humans or other environmental stimuli (e.g., noise)	2
		Signs of a pain face are present and do not disappear in response to the presence of humans or other environmental stimuli (e.g., noise)	4
Head/neck posture (0-2 points)		Head (using the base of the ears as point of reference) is at the same height as the withers or slightly above/below (angle 0 +/- 10°)	0
		Head is held high, clearly (angle > 10°) above the withers	2
		Head is held low, clearly (angle > 10°) below the withers	2
Limb Posture (add all applicable points, 0-6 points)		Weight bearing on all 4 limbs, the horse stands in a rectangle with its forelimbs and hindlimbs parallel to each other and all 4 cannon bones (MCIII and MTIII) visually appr. vertical to the ground (looking at the horse from the side), may rest/cock a hindlimb (may switch rested hindlimb occasionally)	0
		Forelimbs are not parallel to each other (looking at the horse from the side) but the hooves are placed more than one hoof length (10-15cm) apart	2
		Hindlimbs are not parallel to each other (looking at the horse from the side) but the hooves are placed more than two hoof lengths (20-30cm) apart	2
		Both front cannon bones are not vertical (appr. 90°) to the ground, the forelimbs are camped out (placed in front of the vertical) or camped under (placed behind the vertical)	2
		Both rear cannon bones are not vertical (appr. 90°) to the ground, the hindlimbs are camped under (placed in front of the vertical) or camped out (placed behind the vertical)	2
weight shifting (0-4 points)	No weight shifting		0
	Weight shifting from one limb to another (fore- or hindlimbs) 2-4 x/ in 2 minutes (weight shifting by moving hooves or lateral movement of head/neck)		2
	Frequent weight shifting from one limb to another (fore- or hindlimbs) > 4 x/ 2min		4
weight bearing (0-4 points)	Weight bearing on all 4 limbs, may rest/cock a hindlimb (may switch rested hindlimb occasionally)		0
	Repeated lifting of a limb for a few seconds		4
	One limb is only minimally weight bearing or non- weight bearing		4
Lameness at the walk (0-4 points, assess only if the horse is moving without being prompted)	Horse does not move → lameness cannot be assessed		NA
	Normal movement, no lameness evident		0
	Stiff gait, slow, decreased range of motion, no clear lameness, (may be lame when turning)		2
	Clearly lame		4

FIGURE 1
Refined musculoskeletal pain scale.

osteoarthritis ($n = 74/154$, 48%), tendinopathy ($n = 29/154$, 19%), or laminitis ($n = 25/154$, 16%) according to their medical records.

3.2 Horse examination and assessment parameters

The 154 horses had a mean original MPS of 4.8 (SD: 3.0, range: 0–14, median: 4.0, IQR: 2.8–7.0). The 11 horses unable to trot were scored 5 for the subjective lameness exam at the trot and assigned a Q-score of 115 (10% higher than the maximum measured Q-score of 104.7). The mean subjective lameness score (maximum of the four limbs, scale of 0–5) at the walk was 1.6 (SD: 1.2, range: 0–5, median: 2.0, IQR: 0.75–2.0) and at the trot, 2.2 (SD: 1.1, range: 0–5, median: 2.0, IQR: 2.0–3.0). The mean Q-score of the objective lameness exam was 28 mm (SD: 35 mm, range: 0.0–104.7 mm, median: 13 mm, IQR: 8.4–23 mm).

The cohort of 60 horses used to validate the refined MPS had a mean MPS of 6.9 (SD: 4.3, range: 0.0–18.0, median: 6.0, IQR: 4.0–10.0) and a mean lameness of 2.1 at the walk (SD: 1.2, range: 0.0–4.0, median: 2.0, IQR: 1.0–3.0) and 2.7 at the trot (SD: 1.6, range: 0.0–5.0, median: 3.0, IQR: 2.0–3.0). One horse was excluded from the lameness exam owing to chronic ataxia.

3.3 Musculoskeletal pain scale—descriptive statistics

The correlations among items and their contribution to the total score were assessed using correlation analysis and principal component analysis. The correlation analysis using the non-parametric Spearman and the parametric Pearson correlations

showed negligible or positive correlations of varying strengths among the items of the original MPS (Figure 2). In the original MPS, the item “location in the box/enclosure” correlated little with the other items ($r < 0.25$) and had a correspondingly low item–total correlation of 0.092, while the other items correlated with a correlation coefficient between about 0.3–0.4 and had item–total correlations between 0.18 and 0.445. The correlation of the items with the total original MPS is also shown, from which it is evident that location, demeanour, and head–neck posture have a lower correlation than lameness, weight distribution, and pain face, which show pairwise correlations in the range 0.25–0.75. Using the non-parametric Spearman or the parametric Pearson correlation produced qualitatively and quantitatively similar patterns. As expected from the correlation analysis, all subitems, except demeanour and location, contribute positively to the first principal component, especially lameness and pain face, which load highly (Table 1), while they load with opposite signs on the second principal component. All other items generally dominate one further principal component.

3.4 Musculoskeletal pain scale—validity and predictive performance

The MPS correlated significantly with the subjective lameness score at the walk (Spearman $r = 0.51$, $p < 0.0001$) and the trot (Spearman $r = 0.45$, $p < 0.0001$) and the objective lameness measurements (Spearman $r = 0.37$, $p = 0.0001$, $p \leq 0.0001$).

The MPS was significantly different (difference between means: 2.1 ± 0.5 SEM, $p < 0.0001$) between horses that needed veterinary intervention (mean MPS: 6.2, $n = 49$) and those that did not (mean MPS: 4.1, $n = 105$). The MPS was also significantly




FIGURE 2

Pearson correlations for the original (left) and refined (right) MPS for the total and the various sub-items of the MPS. The colours and sizes of the balls indicate the strength of the correlation.

TABLE 1 The proportion of the principal components indicating their relative importance are given in the first row. The loadings of the items, reflecting their contributions to the principal components, are given in the following for the original (A) and refined (B) MPS. Values below 0.1 are not reported, i.e., left blank.

A							
Component	1	2	3	4	5	6	
Location		0.592	0.767	0.243			
Demeanour					0.109	−0.990	
Pain face	0.604	−0.605	0.299	0.404			
Head–neck posture	0.112		0.139	−0.164	−0.963		
Limb posture/weight-bearing/shifting	0.331		0.293	−0.864	0.229		
Lameness	0.712	0.521	−0.465				
Proportion of variance	0.483	0.171	0.159	0.119	0.049	0.019	
B							
Component	1	2	3	4	5	6	7
Demeanour							0.998
Pain face	0.460		−0.747		0.379	0.281	
Head–neck posture	0.127		−0.479	−0.140	−0.670	−0.529	
Limb posture	0.767	0.467	0.390		−0.176		
Weight-bearing	0.204	−0.133	0.108	−0.845	0.371	−0.276	
Weight shifting		−0.166		−0.416	−0.489	0.746	
Lameness	0.371	−0.856	0.221	0.276			
Proportion of variance	0.369	0.218	0.163	0.124	0.0653	0.0549	0.006

different (difference between means: 4.3 ± 0.5 SEM, $p < 0.0001$) between horses that were clearly lame at the walk (lameness ≥ 3 , mean MPS: 8.3, $n = 29$) and those that were not or only mildly lame (grade 0–2, mean MPS: 4.0, $n = 125$). Similarly, the MPS was significantly different (difference between means: 2.9 ± 0.5 SEM, $p < 0.0001$) between horses that were clearly lame at the trot (lameness ≥ 3 , mean MPS: 6.8, $n = 45$) and those that were not or only mildly lame (grade 0–2, mean MPS: 3.9, $n = 109$). Furthermore, the MPS was significantly different between objectively measured lameness scores, specifically between no lameness (Q-score ≤ 8.5 , mean MPS: 3.5) and moderate to severe lameness (Q-score > 30 , mean MPS: 7.9, $p < 0.0001$), and between mild (Q-scores 8.5–30, mean MPS: 4.1) and moderate to severe lameness ($p < 0.0001$), but not between mild and no lameness.

3.5 Musculoskeletal pain scale—interrater agreement and refinement

During the refinement process, the item “location in the box/enclosure” was dropped as it was not consistently possible to reliably rate the location on videos, and the MPS was intended to allow for remote scoring to avoid the confounding effect of rater presence on horses’ behaviour. The original item, scoring limb posture, weight-bearing, and weight shifting together were divided into three items in the refined tool. The definition of the other items was optimized based on feedback from the raters and interrater agreement to optimize clarity and minimize the potential for misinterpretation. The time required to complete pain assessment was reduced from 7 min with the original MPS to 2 min with the refined MPS, hence enhancing its user-friendliness and corresponding clinical and research utility.

Correlations among the items of the refined MPSs (Figure 2) were lower than for the original MPS, indicating lower item redundancy; demeanour correlated little with the other items

($r < 0.25$); pain face correlated moderately with head-neck and limb posture, and weight-bearing correlated with weight shifting, all with a correlation coefficient between 0.3 and 0.4. The correlation of the items with the total refined MPS is below 0.1 for demeanour and between 0.25 and 0.6 for the other items. The item–total correlation for demeanour was only 0.053, that of the rest of the items between 0.19 (weight shifting) and 0.464 (pain face). Using the non-parametric Spearman or the parametric Pearson correlation produced qualitatively and quantitatively similar patterns. The first principal component has positive loadings for all items of the refined score except demeanour and weight shifting (Table 1). While the item demeanour dominates one principal component of the refined MPS, the loadings of all other items are generally more dispersed over the various other components than for the original MPS. The residual variance can be attributed to the variation in the video quality and hence as technical variance.

The intraclass correlation increased from 0.77 [95% confidence interval (CI): 0.62–0.88] for the original MPS to 0.88 (95% CI: 0.80–0.94) for the refined MPS tool (Supplementary Table S1).

A mixed model analysis evaluated the variance due to variability among horses, among raters, and the residual variance (Table 2). The variability among horses is more than five times higher than that among raters for the total scores as well as

TABLE 2 Mixed model analysis of the total refined MPS and its various subitems.

Variance component	Horse-ID	Rater-ID	Residual
Refined MPS	6.935	0.820	5.313
Demeanour	0.020	0.003	0.158
Pain face	0.555	0.004	1.228
Head-neck posture	0.301	0.003	0.502
Limb posture	1.116	0.482	1.260
Weight-bearing	0.771	0.331	1.429
Weight shifting	0.523	0.000	0.333
Lameness	-	-	-

lameness, head-neck posture, and limb posture, indicating that these variables can be reliably scored by the different raters. The ratio is less favourable for pain face, weight-bearing, and weight shifting. The residual variance can be attributed to the variation in the video quality and hence as technical variance.

3.6 Refined musculoskeletal pain scale—validity and discriminative power

The refined MPS was validated by its significant correlation with the subjective lameness score in walk (Spearman $r = 0.44$, $p = 0.001$) and trot ($r = 0.5$, $p < 0.0001$).

The MPS was significantly different (difference between means: 3.8 ± 1.1 SEM, $p = 0.0009$) between horses that were clearly lame in

walk (lameness ≥ 3 , mean MPS: 9.3, $n = 21$) and those that were not or only mildly lame (grade 0–2, mean MPS: 5.6, $n = 38$, Figure 3). Furthermore, the MPS was significantly different (difference between means: 4.2 ± 0.99 SEM, $p < 0.0001$) between horses that were clearly lame at the trot (lameness ≥ 3 , mean MPS: 8.9, $n = 31$) and those that were not or only mildly lame (grade 0–2, mean MPS: 4.7, $n = 28$, Figure 3).

The refined MPS was also significantly different (difference between means: 3.6 ± 1.0 SEM, $p = 0.0007$) between horses that needed veterinary intervention (mean MPS: 8.6, $n = 31$) and those that did not (mean MPS: 5.0, $n = 29$, Figure 4), hence establishing the predictive performance of the MPS for longitudinal health outcomes.

Lastly, the MPS was significantly different (difference between means: 3.2 ± 1.093 SEM, $p = 0.0168$) in horses receiving analgesia between before and after treatment, confirming its responsiveness (Figure 4).

ROC analysis showed the refined MPS to be moderately accurate in discriminating between horses that needed veterinary intervention and those that did not, with a c-statistic of 0.74 (standard error: 0.064, $p = 0.0002$). The ROC analysis yielded several possible MPS cut-off values with corresponding trade-offs in sensitivity and specificity (Table 3, Figure 4, Supplementary Table S2), with an MPS cut-off > 8 providing the best overall combination of sensitivity (51.61%, 95% CI: 34.84%–68.03%) and specificity (89.66%, 95% CI: 73.61%–96.42%) with a Youden index (=specificity plus sensitivity minus one) of 0.41, indicating that horses with an MPS of 8 or greater have a high probability of a painful condition and therefore should be further examined.

Without the item lameness, ROC analysis (c-statistic: 0.65, standard error: 0.071, $p = 0.034$) showed that an MPS cut-off of > 4 (Youden index of 0.264) yielded the best combination of sensitivity (67.74%, 95% CI: 51.6%–74.2%) and specificity (58.62%, 95% CI: 48.4%–75.5%, Supplementary Figure S2).





TABLE 3 Sensitivity and specificity plus 95% CI calculated by ROC analysis for each MPS cut-off value for identifying horses needing veterinary intervention.

Cut-off	Sensitivity (%)	95% CI	Specificity (%)	95% CI	Youden index	Likelihood ratio
Differentiation between horses that needed medical intervention and those that did not						
>0	96.77	83.81%–99.83%	13.79	5.5%–30.56%	0.11	1.123
>2.0	90.37	75.10%–96.65%	31.03	17.28%–49.23%	0.21	1.31
>4.0	77.42	60.19%–88.60%	51.72	34.43%–68.61%	0.29	1.6
>6.0	61.29	43.82%–76.27%	68.97	50.77%–82.72%	0.3	1.975
>8.0	51.61	34.84%–68.03%	89.66	73.61%–96.42%	0.41	4.989
>10.0	32.26	18.57%–49.86%	93.1	78.04%–98.77%	0.25	4.677
>12.0	16.13	7.093%–32.63%	100.0	88.30%–100.0%	0.16	
>16.0	3.226	0.166%–16.19%	100.0	88.30%–100.0%	0.03	

The best Youden index (the sum of sensitivity and specificity – 1) is indicated in bold.

The difference in the MPSs (difference between means: 2.0 ± 0.9) between horses that needed veterinary intervention (mean: 6.3) and those that did not (mean: 4.3) remained statistically significant also without the lameness item ($p = 0.0331$, Supplementary Figure S2).

4 Discussion

The high prevalence and impact of chronic musculoskeletal conditions and the poor recognition of lameness and the associated pain necessitate the inclusion of pain as a fourth vital sign in the routine evaluation of all horses to facilitate appropriate treatment and improve equine welfare (52, 86, 87). Regular pain assessment using a reliable, valid, and clinically useful tool would enable the identification of pain, timely interventions, monitoring treatment effects, and facilitate communication among veterinarians and caretakers. This study established, refined, and validated a multidimensional MPS and demonstrated its predictive performance for longitudinal health outcomes, discriminative power, and good interrater agreement between veterinary practitioners and equestrian raters. The MPS was validated as a measure for equestrians and veterinarians alike to assess horses for the presence of painful conditions and monitor the efficacy of treatment interventions, not just in a hospital setting but especially also in their home environment. The MPS is based on several pain behaviours, including features of the equine pain face, postural indicators, demeanour, and lameness to accommodate for individual variations in pain behaviour and differences in pain behaviours between acute and chronic pain (13, 26, 40, 41, 88–91).

Pain is a complex, uniquely individual, unpleasant experience associated with actual or potential tissue damage, encompassing both sensory (intensity) and affective (unpleasantness) components (92). The affective dimension of pain is associated with behavioural changes aimed at avoiding pain and minimizing injury (23, 54, 93). While acute pain is a protective response to noxious stimuli, chronic pain persists beyond the expected healing time and may be either a symptom of chronic peripheral disease, maladaptive nervous system dysfunction, or both (89, 90, 94–100). Acute pain tends to respond to anti-inflammatory pain relief. Decreasing the load on the affected area by postural adaptations and guarding during movement and at rest may reduce acute pain

(13, 40, 41). By contrast, in chronic pain, due to the central sensitization that is often present, the degree, duration, and spatial extent of pain may be increased and distorted, leading to more widespread pain and multisite hyperalgesia and allodynia (13, 40, 86, 88, 89, 95, 101–103). Therefore, once central sensitization has occurred, pain perception may no longer reflect the presence, intensity, or duration of peripheral noxious inputs (103, 104). Accordingly, behavioural changes associated with chronic pain can vary greatly, necessitating a multidimensional pain assessment tool that includes the effect of pain on demeanour, functional assessments (lameness), and different pain behaviours (24, 41, 86, 89–91, 94–97, 99, 105, 106). Pain behaviours can be categorized into two overlapping groups: protective and communicative pain behaviours (106). Protective pain behaviours, such as postural adjustments and guarding, are often directly associated with the painful area (13, 40). By contrast, communicative pain behaviours, including facial expressions, are universal indicators of pain and mechanisms for communicating pain to conspecifics (13, 40).

Posture, the dynamic alignment and positioning of the body orchestrated by the neuromuscular system, is often mistakenly conflated with conformation, which pertains to the static skeletal architecture and body proportions (13, 33, 40, 45, 47, 48, 107, 108), confounding research into the complex interplay between posture and musculoskeletal pain. Posture facilitates efficient weight distribution across the musculoskeletal system, balancing the centre of gravity over the base of support to minimize energy expenditure and stress on anatomical structures (33, 109, 110). As symmetrical loading of the limbs provides the greatest biomechanical stability and hence requires the least corrective actions and energy to maintain balance (110), sound horses exhibit a symmetrical weight distribution, with approximately 60% of the weight borne by the forelimbs and 40%, by the hindlimbs (33, 111, 112). Conversely, horses afflicted with orthopaedic conditions may attempt to alleviate pain by shifting the weight away from the affected limb, effectively altering their centre of gravity (113). Weight-bearing and stance asymmetry may therefore signal pain relieved by adopting this posture (112). Similarly, an elevated neck posture has been identified as a potential indicator of underlying back disorders in horses (53, 56, 57, 107, 114, 115). Therefore, the MPS includes head-neck posture, limb posture, weight-bearing, and weight shifting as separate items to reflect the postural adaptations commonly

observed in response to orthopaedic pain. We note that due to the complexity of pain, a single item may not correlate highly with other items or show a low item–total correlation, i.e., pain may represent more than one dimension. This may explain why the factor loadings are relatively low and the proportion of the variance explained by the principle components is rather even, especially with the refined MPS. In particular, pain face may integrate many aspects of pain, while other items may reflect a specific condition or individualized reaction.

The horses' interaction with humans may also be variably affected by pain (25, 38, 44). Depending on the intensity of the noxious stimuli and the familiarity of the environment and observer, painful horses may either be reluctant to interact with humans or increase their contact-seeking behaviour (25, 38, 44). Conversely, the horses may reduce or relieve pain behaviour when people approach or interact (46), which can lead to underestimation of the pain and subsequent therapeutic deficits and welfare problems. Therefore, the MPS was designed to be applicable from a distance to avoid disrupting pain behaviours.

This study established the content, criterion, and construct validity of the MPS using an expert panel for content validation, correlation with lameness for criterion validation, and correlation with longitudinal health outcomes for construct validation. As criterion validation assesses how accurately a scale reflects the *gold standard* for measuring the same construct (63), the lacking *gold standard* or other previously validated method for measuring the individual experience of pain is one of the main limitations of this study, which uses lameness as an indicator of orthopaedic pain. Although lameness is a reliable indicator of pain, the absence of overt lameness does not exclude the possibility of pain. This limitation, the lacking gold standard and objective, of quantitative pain measurement also extends to the evaluation of the MPS's construct validity, the assessment of its ability to discriminate between horses in pain and pain-free horses (63). This study uses the need for veterinary intervention, identified by the staff of the sanctuary, to assess the MPS's construct validity. The inherent subjectivity of this assessment is however mitigated by the horses statistically significant reduction in pain in response to analgesia. However, multicentre studies using larger patient cohorts are needed to further evaluate the MPS's utility to identify horses in pain in various husbandry and demographic settings.

While the MPS is a quantitative tool, it is crucial to recognize that pain expression does not directly correlate with the severity of tissue damage but reflects horses' individual experience and personality and that many pain behaviours are part of the communication repertoire of healthy horses as well (26). However, the MPS correlates well with lameness at the walk and trot and showed very good predictive performance for longitudinal health outcomes and discriminative power in identifying lame horses and horses needing veterinary intervention. In this cohort, the maximum MPS was 18 (the horse was 3/5 lame at the walk and too lame to trot), the maximum score of 26 was not reached by any horse, possibly because no horse in this study suffered from severe pain. Based on the ROC analysis and the differences between horses

exhibiting obvious lameness (≥ 3 on a scale of 0–5) or requiring veterinary intervention, as opposed to horses with minimal or no observable health concerns, horses with an MPS exceeding 8 (or 4 if the lameness item cannot be assessed) should undergo further examination to identify the underlying cause and determine if treatment is necessary.

5 Conclusions

In summary, the MPS showed good interrater repeatability between expert and lay scorers, significant correlation with lameness at the walk and trot, and good predictive validity for longitudinal health outcomes, confirming its ability to identify horses with musculoskeletal pain. Given the prevalence of chronic musculoskeletal conditions, the poor recognition of lameness, and the suffering caused by unrelieved pain, pain assessment should be included in all veterinary examinations, and caretakers should regularly evaluate their horse's pain status to facilitate timely therapeutic interventions. Routine pain assessment using a reliable and validated tool may help address the widespread problem of unrelieved pain already voiced by the philosopher Michel de Montaigne in 1589: "For heaven's sake, let medicine someday give me some good and perceptible relief and you will see how I shall cry out in good earnest: At last I yield to an efficient science."

Data availability statement

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

Ethics statement

The requirement of ethical approval was waived by the Institutional Ethics Committee of the University of Veterinary Medicine, Vienna, for the studies involving animals because this study was non-invasive and entailed only monitoring the horses under their current conditions of life. No specific veterinary treatments or interventions were carried out for the purpose of this study. The studies were conducted in accordance with the local legislation and institutional requirements.

Author contributions

UA: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Validation, Visualization, Writing – original draft, Writing – review & editing. ZK: Data curation, Investigation, Project administration, Writing – review & editing. CV: Formal analysis, Methodology, Software, Validation, Visualization, Writing – review & editing. SR: Investigation, Writing – review & editing. RH: Investigation, Writing – review & editing. LT: Investigation, Writing – review & editing. CG: Investigation, Writing – review & editing. JB: Investigation,

Writing – review & editing. FJ: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Validation, Visualization, Writing – original draft, Writing – review & editing.

Funding

The authors declare that no financial support was received for the research, authorship, and/or publication of this article.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

- Egenvall A, Lönnell C, Roepstorff L. Analysis of morbidity and mortality data in riding school horses, with special regard to locomotor problems. *Prev Vet Med.* (2009) 88:193–204. doi: 10.1016/j.prevetmed.2008.10.004
- Weeren van PR, Back W. Musculoskeletal disease in aged horses and its management. *Vet Clin North Am Equine Pract.* (2016) 32:229–47. doi: 10.1016/j.cveq.2016.04.003
- Ireland JL, Clegg PD, McGowan CM, McKane SA, Chandler KJ, Pinchbeck GL. Disease prevalence in geriatric horses in the United Kingdom: veterinary clinical assessment of 200 cases. *Equine Vet J.* (2012) 44:101–6. doi: 10.1111/j.2042-3306.2010.00361.x
- Ireland JL, McGowan CM, Clegg PD, Chandler KJ, Pinchbeck GL. A survey of health care and disease in geriatric horses aged 30 years or older. *Vet J.* (2012) 192:57–64. doi: 10.1016/j.tvjl.2011.03.021
- Dyson S, Berger J, Ellis AD, Mullard J. Development of an ethogram for a pain scoring system in ridden horses and its application to determine the presence of musculoskeletal pain. *J Vet Behav.* (2018) 23:47–57. doi: 10.1016/j.jveb.2017.10.008
- Dunn KM, Campbell P, Lewis M, Hill JC, Windt van der DA, Afolabi E, et al. Refinement and validation of a tool for stratifying patients with musculoskeletal pain. *Eur J Pain.* (2021) 25:2081–93. doi: 10.1002/ejp.1821
- Safiri S, Kolahi A, Cross M, Hill C, Smith E, Carson-Chahhoud K, et al. Prevalence, deaths, and disability-adjusted life years due to musculoskeletal disorders for 195 countries and territories 1990–2017. *Arthritis Rheumatol.* (2021) 73:702–14. doi: 10.1002/art.41571
- Woolf AD. Global burden of osteoarthritis and musculoskeletal diseases. *BMC Musculoskelet Disord.* (2015) 16:S3. doi: 10.1186/1471-2474-16-s1-s3
- Egenvall A, Bonnett B, Wattle O, Emanuelson U. Veterinary-care events and costs over a 5-year follow-up period for warmblooded riding horses with or without previously recorded locomotor problems in Sweden. *Prev Vet Med.* (2008) 83:130–43. doi: 10.1016/j.prevetmed.2007.06.008
- Ireland JL, Clegg PD, McGowan CM, McKane SA, Pinchbeck GL. A cross-sectional study of geriatric horses in the United Kingdom. Part 2: health care and disease. *Equine Vet J.* (2010) 43:37–44. doi: 10.1111/j.2042-3306.2010.00142.x
- Ireland JL, Clegg PD, McGowan CM, McKane SA, Chandler KJ, Pinchbeck GL. Comparison of owner-reported health problems with veterinary assessment of geriatric horses in the United Kingdom. *Equine Vet J.* (2011) 44:94–100. doi: 10.1111/j.2042-3306.2011.00394.x
- Jönsson L, Roepstorff L, Egenvall A, Näsholm A, Dalin G, Philipsson J. Prevalence of clinical findings at examinations of young Swedish warmblood riding horses. *Acta Vet Scand.* (2013) 55:34. doi: 10.1186/1751-0147-55-34
- Ask K, Andersen PH, Tamminen L-M, Rhodin M, Hernlund E. Performance of four equine pain scales and their association to movement asymmetry in horses with induced orthopedic pain. *Frontiers Vet Sci.* (2022) 9:938022. doi: 10.3389/fvets.2022.938022
- Lillie AK, Read S, Mallen C, Croft P, McBeth J. Musculoskeletal pain in older adults at the end-of-life: a systematic search and critical review of the literature with priorities for future research. *BMC Palliat Care.* (2013) 12:27–27. doi: 10.1186/1472-684x-12-27
- Gillsjö C, Nässén K, Berglund M. Suffering in silence: a qualitative study of older adults' experiences of living with long-term musculoskeletal pain at home. *Eur J Ageing.* (2021) 18:55–63. doi: 10.1007/s10433-020-00566-7
- Levetown M, Reid MC. What is the role and impact of osteoarthritis in the realm of palliative care? *J Palliative Care.* (2014) 30:317–20. doi: 10.1177/082585971403000415
- Rhodin M, Egenvall A, Andersen PH, Pfau T. Head and pelvic movement asymmetries at trot in riding horses in training and perceived as free from lameness by the owner. *PLoS One.* (2016) 12:e0176253. doi: 10.1371/journal.pone.0176253
- Müller-Quirin J, Dittmann MT, Roepstorff C, Arpagaus S, Latif SN, Weishaupt MA. Riding soundness—comparison of subjective with objective lameness assessments of owner-sound horses at trot on a treadmill. *J Equine Vet Sci.* (2020) 95:103314. doi: 10.1016/j.jevs.2020.103314
- Welsh TP, Yang AE, Makris UE. Musculoskeletal pain in older adults: a clinical review. *Med Clin N Am.* (2020) 104:855–72. doi: 10.1016/j.mcna.2020.05.002
- Cowen R, Stasiowska MK, Laycock H, Bantel C. Assessing pain objectively: the use of physiological markers. *Anaesthesia.* (2015) 70:828–47. doi: 10.1111/anae.13018
- Arendt-Nielsen L. Evaluation of pain in humans. In: Gebhart GF, Schmidt RF, editors. *Encyclopedia of pain.* Springer, Berlin, Heidelberg, Germany (2013). p. 1204–16. doi: 10.1007/978-3-642-28753-4_1387
- Carvalho JRG, Trindade PHE, Conde G, Antonioli ML, Funnicelli MIG, Dias PP, et al. Facial expressions of horses using weighted multivariate statistics for assessment of subtle local pain induced by polylactide-based polymers implanted subcutaneously. *Animals.* (2022) 12:2400. doi: 10.3390/ani12182400
- Ashley FH, Waterman-Pearson AE, Whay HR. Behavioural assessment of pain in horses and donkeys: application to clinical practice and future studies. *Equine Vet J.* (2005) 37:565–75. doi: 10.2746/042516405775314826
- Kunz M, Lautenbacher S, LeBlanc N, Rainville P. Are both the sensory and the affective dimensions of pain encoded in the face? *Pain.* (2012) 153:350–8. doi: 10.1016/j.pain.2011.10.027
- Price J, Catriona S, Welsh EM, Waran NK. Preliminary evaluation of a behaviour-based system for assessment of post-operative pain in horses following arthroscopic surgery. *Vet Anaesth Analg.* (2003) 30:124–37. doi: 10.1046/j.1467-2995.2003.00139.x
- Ijichi C, Collins LM, Elwood RW. Pain expression is linked to personality in horses. *Appl Anim Behav Sci.* (2014) 152:38–43. doi: 10.1016/j.applanim.2013.12.007
- Costa ED, Minero M, Lebelt D, Stucke D, Canali E, Leach MC. Development of the horse grimace scale (HGS) as a pain assessment tool in horses undergoing routine castration. *PLoS One.* (2014) 9:e92281. doi: 10.1371/journal.pone.0092281
- Arbour C, Gélinas C. Are vital signs valid indicators for the assessment of pain in postoperative cardiac surgery ICU adults? *Intensiv Crit Care Nurs.* (2010) 26:83–90. doi: 10.1016/j.jccn.2009.11.003
- Gélinas C, Arbour C. Behavioral and physiologic indicators during a nociceptive procedure in conscious and unconscious mechanically ventilated adults: similar or different? *J Crit Care.* (2009) 24:628.e7–17. doi: 10.1016/j.jccr.2009.01.013
- Visser E. What is pain? II: pain expression and behaviour, evolutionary concepts, models and philosophies. *Australian Anaesthesia.* (2009) 31:35–43.
- Alamam DM, Leaver A, Moloney N, Alsobayel HI, Alashaikh G, Mackey MG. Pain behaviour scale (PaBS): an exploratory study of reliability and construct

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpain.2023.1292299/full#supplementary-material>

validity in a chronic low back pain population. *Pain Res Manag.* (2019) 2019:2508019. doi: 10.1155/2019/2508019

32. Main CJ, Watson PJ. Guarded movements. *J Musculoskelet Pain.* (2010) 4:163–70. doi: 10.1300/j094v04n04_16

33. Gellman K, Ruina A. Standing horse posture: a longer stance is more stable. *Biol Open.* (2022) 11:bio059139. doi: 10.1242/bio.059139

34. Cramer H, Mehling WE, Saha FJ, Dobos G, Lauche R. Postural awareness and its relation to pain: validation of an innovative instrument measuring awareness of body posture in patients with chronic pain. *BMC Musculoskelet Disord.* (2018) 19:109. doi: 10.1186/s12891-018-2031-9

35. Walsh J, Eccleston C, Keogh E. Pain communication through body posture: the development and validation of a stimulus set. *Pain.* (2014) 155:2282–90. doi: 10.1016/j.pain.2014.08.019

36. Werner P, Al-Hamadi A, Limbrecht-Ecklundt K, Walter S, Traue HC. Head movements and postures as pain behavior. *PLoS One.* (2018) 13:e0192767. doi: 10.1371/journal.pone.0192767

37. McLennan KM, Rebelo CJB, Corke MJ, Holmes MA, Leach MC, Constantino-Casas F. Development of a facial expression scale using footrot and mastitis as models of pain in sheep. *Appl Anim Behav Sci.* (2016) 176:19–26. doi: 10.1016/j.applanim.2016.01.007

38. Gleerup KB, Forkman B, Lindegaard C, Andersen PH. An equine pain face. *Vet Anaesth Analg.* (2015) 42:103–14. doi: 10.1111/vaa.12212

39. Aviezer H, Trope Y, Todorov A. Body cues, not facial expressions, discriminate between intense positive and negative emotions. *Science.* (2012) 338:1225–9. doi: 10.1126/science.1224313

40. Ask K, Rhodin M, Tamminen L-M, Hernlund E, Andersen PH. Identification of body behaviors and facial expressions associated with induced orthopedic pain in four equine pain scales. *Animals (Basel).* (2020) 10:2155. doi: 10.3390/ani10112155

41. Prkachin KM, Schultz IZ, Hughes E. Pain behavior and the development of pain-related disability: the importance of guarding. *Clin J Pain.* (2007) 23:270–7. doi: 10.1097/ajp.0b013e3180308d28

42. Prkachin KM, Berzins S, Mercer SR. Encoding and decoding of pain expressions: a judgement study. *Pain.* (1994) 58:253–9. doi: 10.1016/0304-3959(94)90206-2

43. LeResche L, Dworkin SF. Facial expression accompanying pain. *Soc Sci Med.* (1984) 19:1325–30. doi: 10.1016/0277-9536(84)90020-0

44. Bussi eres G, Jacques C, Lainay O, Beauchamp G, Leblond A, Cadore J-L, et al. Development of a composite orthopaedic pain scale in horses. *Res Vet Sci.* (2008) 85:294–306. doi: 10.1016/j.rvsc.2007.10.011

45. Torcivia C, McDonnell S. Equine discomfort ethogram. *Animals (Basel).* (2021) 11:580. doi: 10.3390/ani11020580

46. Torcivia C, McDonnell S. In-Person caretaker visits disrupt ongoing discomfort behavior in hospitalized equine orthopedic surgical patients. *Animals (Basel).* (2020) 10:210. doi: 10.3390/ani10020210

47. Loon van JPAM, Macri L. Objective assessment of chronic pain in horses using the horse chronic pain scale (HCPS): a scale-construction study. *Animals (Basel).* (2021) 11:1826. doi: 10.3390/ani11061826

48. Loon van JPAM, Dierendonck MCV. Pain assessment in horses after orthopaedic surgery and with orthopaedic trauma. *Vet J.* (2019) 246:85–91. doi: 10.1016/j.tvjl.2019.02.001

49. Price J, Clarke N, Welsh EM, Waran N. Preliminary evaluation of subjective scoring systems for assessment of postoperative pain in horses. *Vet Anaesth Analg.* (2003) 30:97. doi: 10.1046/j.1467-2995.2003.00132_15.x

50. Gleerup KB, Lindegaard C. Recognition and quantification of pain in horses: a tutorial review. *Equine Vet Educ.* (2015) 28:47–57. doi: 10.1111/eve.12383

51. Hug F, Hodges PW, Salomoni SE, Tucker K. Insight into motor adaptation to pain from between-leg compensation. *Eur J Appl Physiol.* (2014) 114:1057–65. doi: 10.1007/s00421-014-2840-y

52. Monteiro BP, Lascelles BDX, Murrell J, Robertson S, Steagall PVM, Wright B. 2022 WSAVA guidelines for the recognition, assessment and treatment of pain. *J Small Anim Pr.* (2023) 64:177–254. doi: 10.1111/jsap.13566

53. S  n  que E, Lesimple C, Morisset S, Hausberger M. Could posture reflect welfare state? A study using geometric morphometrics in riding school horses. *PLoS One.* (2019) 14:e0211852. doi: 10.1371/journal.pone.0211852

54. Broom   S, Ask K, Rashid-Engstr  m M, Andersen PH, Kjellstr  m H. Sharing pain: using pain domain transfer for video recognition of low grade orthopedic pain in horses. *PLoS One.* (2022) 17:e0263854. doi: 10.1371/journal.pone.0263854

55. Dyson S, Berger JM, Ellis AD, Mullard J. Behavioral observations and comparisons of nonlame horses and lame horses before and after resolution of lameness by diagnostic analgesia. *J Vet Behav.* (2018) 26:64–70. doi: 10.1016/j.jveb.2018.05.001

56. S  n  que E, Morisset S, Lesimple C, Hausberger M. Testing optimal methods to compare horse postures using geometric morphometrics. *PLoS One.* (2018) 13:e0204208. doi: 10.1371/journal.pone.0204208

57. Lesimple C, Fureix C, Margerie ED, S  n  que E, Menguy H, Hausberger M. Towards a postural indicator of back pain in horses (*Equus caballus*). *PLoS One.* (2012) 7:e44604. doi: 10.1371/journal.pone.0044604

58. Mokkink LB, Boers M, Vleuten van der CPM, Bouter LM, Alonso J, Patrick DL, et al. Cosmin risk of bias tool to assess the quality of studies on reliability or measurement error of outcome measurement instruments: a Delphi study. *BMC Med Res Methodol.* (2020) 20:293. doi: 10.1186/s12874-020-01179-5

59. Mokkink LB, Terwee CB, Gibbons E, Stratford PW, Alonso J, Patrick DL, et al. Inter-rater agreement and reliability of the COSMIN (Consensus-based standards for the selection of health status measurement instruments) checklist. *BMC Med Res Methodol.* (2010) 10:82. doi: 10.1186/1471-2288-10-82

60. Mokkink LB, Terwee CB, Patrick DL, Alonso J, Stratford PW, Knol DL, et al. The COSMIN checklist for assessing the methodological quality of studies on measurement properties of health status measurement instruments: an international Delphi study. *Qual Life Res.* (2010) 19:539–49. doi: 10.1007/s11136-010-9606-8

61. Mokkink LB, Terwee CB, Knol DL, Stratford PW, Alonso J, Patrick DL, et al. The COSMIN checklist for evaluating the methodological quality of studies on measurement properties: a clarification of its content. *BMC Med Res Methodol.* (2010) 10:22. doi: 10.1186/1471-2288-10-22

62. Mokkink LB, Terwee CB, Patrick DL, Alonso J, Stratford PW, Knol DL, et al. The COSMIN study reached international consensus on taxonomy, terminology, and definitions of measurement properties for health-related patient-reported outcomes. *J Clin Epidemiology.* (2010) 63:737–45. doi: 10.1016/j.jclinepi.2010.02.006

63. Tomacheuski RM, Monteiro BP, Evangelista MC, Luna SPL, Steagall PV. Measurement properties of pain scoring instruments in farm animals: a systematic review using the COSMIN checklist. *PLoS One.* (2023) 18:e0280830. doi: 10.1371/journal.pone.0280830

64. Streiner DL, Norman GR, Fulton C. Health measurement scales: a practical guide to their development and use. *Int J Rehabilitation Res.* (1991) 14:364. doi: 10.1097/00004356-199112000-00017

65. Sutton GA, Dahan R, Turner D, Paltiel O. A behaviour-based pain scale for horses with acute colic: scale construction. *Vet J.* (2013) 196:394–401. doi: 10.1016/j.tvjl.2012.10.008

66. Pritchett LC, Ulibarri C, Roberts MC, Schneider RK, Sellon DC. Identification of potential physiological and behavioral indicators of postoperative pain in horses after exploratory celiotomy for colic. *Appl Anim Behav Sci.* (2003) 80:31–43. doi: 10.1016/s0168-1591(02)00205-8

67. Loon van JPAM, Dierendonck MCV. Monitoring acute equine visceral pain with the equine Utrecht University scale for composite pain assessment (EQUUS-COMPASS) and the equine Utrecht university scale for facial assessment of pain (EQUUS-FAP): a scale-construction study. *Vet J.* (2015) 206:356–64. doi: 10.1016/j.tvjl.2015.08.023

68. Andersen PH, Broom   S, Rashid M, Lundblad J, Ask K, Li Z, et al. Towards machine recognition of facial expressions of pain in horses. *Animals (Basel).* (2021) 11:1643. doi: 10.3390/ani11061643

69. Costa ED, Stucke D, Dai F, Minero M, Leach M, Lebelt D. Using the horse grimace scale (HGS) to assess pain associated with acute laminitis in horses (*Equus caballus*). *Animals (Basel).* (2016) 6:47. doi: 10.3390/ani6080047

70. Schieder K, Zsoldos RR, Dippel M, Siedler C, Tichy A, Licka TF. Use of physical self-experience for teaching lameness evaluation: short-term effects on lameness evaluation of horses with mild forelimb lameness by novice veterinary students. *J Vet Med Educ.* (2020) 47:342–55. doi: 10.3138/jyme.0618-079r

71. Keegan KG. Evidence-based lameness detection and quantification. *Vet Clin North Am Equine Pract.* (2007) 23:403–23. doi: 10.1016/j.cveq.2007.04.008

72. Keegan KG, Yonezawa Y, Pai PF, Wilson DA, Kramer J. Evaluation of a sensor-based system of motion analysis for detection and quantification of forelimb and hind limb lameness in horses. *Am J Vet Res.* (2004) 65:665–70. doi: 10.2460/ajvr.2004.65.665

73. Keegan KG, Yonezawa Y, Pai PF, Wilson DA. Accelerometer-based system for the detection of lameness in horses. *Biomed Sci Instrum.* (2002) 38:107–12.

74. Donnell JR, Frisbie DD, King MR, Goodrich LR, Haussler KK. Comparison of subjective lameness evaluation, force platforms and an inertial-sensor system to identify mild lameness in an equine osteoarthritis model. *Vet J.* (2015) 206:136–42. doi: 10.1016/j.tvjl.2015.08.004

75. Keegan KG, Kramer J, Yonezawa Y, Maki H, Pai PF, Dent EV, et al. Assessment of repeatability of a wireless, inertial sensor-based lameness evaluation system for horses. *Am J Vet Res.* (2011) 72:1156–63. doi: 10.2460/ajvr.72.9.1156

76. Kramer J, Keegan KG, Kelmer G, Wilson DA. Objective determination of pelvic movement during hind limb lameness by use of a signal decomposition method and pelvic height differences. *Am J Vet Res.* (2004) 65:741–7. doi: 10.2460/ajvr.2004.65.741

77. Sk  ldebrand E, Adepu S, L  tzelschwab C, Nystr  m S, Lindahl A, Abrahamsson-Aurell K, et al. A randomized, triple-blinded controlled clinical study with a novel disease-modifying drug combination in equine lameness-associated osteoarthritis. *Osteoarthr Cartil Open.* (2023) 5:100381. doi: 10.1016/j.ocarto.2023.100381

78. Grady SE, Lescun TB, Moore GE, Cooper BR, Davern AJ, Brunner TJ, et al. Ketorolac is not more effective than flunixin meglumine or phenylbutazone in reducing foot pain in horses. *J Equine Vet Sci.* (2020) 94:103204. doi: 10.1016/j.jevs.2020.103204
79. Taschetto PM, Azevedo da MS, Rodrigues da APC, Martini LG, Siqueira DFC, Bernardes AS, et al. Objective lameness assessment in horses used for equine-assisted therapy in Rio Grande do Sul State, Brazil. *Ciência Rural.* (2021) 52:e20200185. doi: 10.1590/0103-8478cr20200185
80. Hardeman AM, Egenvall A, Bragança FMS, Swagemakers J, Koene MHW, Roepstorff L, et al. Visual lameness assessment in comparison to quantitative gait analysis data in horses. *Equine Vet J.* (2022) 54:1076–85. doi: 10.1111/evj.13545
81. Altman DG. *Practical statistics for medical research. 1st ed.* New York: Chapman and Hall/CRC (1990). doi: 10.1201/9780429258589
82. Eisenmann JC, Laurson KR, DuBose KD, Smith BK, Donnelly JE. Construct validity of a continuous metabolic syndrome score in children. *Diabetol Metab Syndr.* (2010) 2:8. doi: 10.1186/1758-5996-2-8
83. Swets JA. Measuring the accuracy of diagnostic systems. *Science.* (1988) 240:1285–93. doi: 10.1126/science.3287615
84. Greiner M, Pfeiffer D, Smith RD. Principles and practical application of the receiver-operating characteristic analysis for diagnostic tests. *Prev Vet Med.* (2000) 45:23–41. doi: 10.1016/s0167-5877(00)00115-x
85. Team RC. *R: A Language and Environment for Statistical Computing.* Vienna: R Foundation for Statistical Computing (2017). Available online at: <https://www.R-project.org> (accessed March 14, 2023).
86. Tompkins DA, Hobelmann JG, Compton P. Providing chronic pain management in the “fifth vital sign” era: historical and treatment perspectives on a modern-day medical dilemma. *Drug Alcohol Depend.* (2017) 173:S11–21. doi: 10.1016/j.drugalcdep.2016.12.002
87. Max MB, Donovan M, Miaskowski CA, Ward SE, Gordon D, Bookbinder M, et al. Quality improvement guidelines for the treatment of acute pain and cancer pain. *JAMA.* (1995) 274:1874–80. doi: 10.1001/jama.1995.03530230060032
88. Cohen SP, Vase L, Hooten WM. Chronic pain: an update on burden, best practices, and new advances. *Lancet.* (2021) 397:2082–97. doi: 10.1016/s0140-6736(21)00393-7
89. Puntillo F, Giglio M, Paladini A, Perchiazzi G, Viswanath O, Urits I, et al. Pathophysiology of musculoskeletal pain: a narrative review. *Ther Adv Musculoskelet Dis.* (2021) 13:1759720X21995067. doi: 10.1177/1759720(21995067)
90. Groh A, Krieger P, Mease RA, Henderson L. Acute and chronic pain processing in the thalamocortical system of humans and animal models. *Neuroscience.* (2018) 387:58–71. doi: 10.1016/j.neuroscience.2017.09.042
91. Fureix C, Menguy H, Hausberger M. Partners with bad temper: reject or cure? A study of chronic pain and aggression in horses. *PLoS One.* (2010) 5:e12434. doi: 10.1371/journal.pone.0012434
92. Talbot K, Madden VJ, Jones SL, Moseley GL. The sensory and affective components of pain: are they differentially modifiable dimensions or inseparable aspects of a unitary experience? A systematic review. *Br J Anaesth.* (2019) 123:e263–72. doi: 10.1016/j.bja.2019.03.033
93. Sneddon LU, Elwood RW, Adamo SA, Leach MC. Defining and assessing animal pain. *Anim Behav.* (2014) 97:201–12. doi: 10.1016/j.anbehav.2014.09.007
94. McLennan KM, Miller AL, Costa ED, Stucke D, Corke MJ, Broom DM, et al. Conceptual and methodological issues relating to pain assessment in mammals: the development and utilisation of pain facial expression scales. *Appl Anim Behav Sci.* (2019) 217:1–15. doi: 10.1016/j.applanim.2019.06.001
95. Woller SA, Eddinger KA, Corr M, Yaksh TL. An overview of pathways encoding nociception. *Clin Exp Rheumatol.* (2017) 35(Suppl 107):40–6.
96. Vardeh D, Mannion RJ, Woolf CJ. Toward a mechanism-based approach to pain diagnosis. *J Pain.* (2016) 17:T50–69. doi: 10.1016/j.jpain.2016.03.001
97. Gold MS, Gebhart GF. Nociceptor sensitization in pain pathogenesis. *Nat Med.* (2010) 16:1248–57. doi: 10.1038/nm.2235
98. Fornasari D. Pain mechanisms in patients with chronic pain. *Clin Drug Invest.* (2012) 32(Suppl 1):45–52. doi: 10.2165/11630070-000000000-00000
99. Basbaum AI, Bautista DM, Scherrer G, Julius D. Cellular and molecular mechanisms of pain. *Cell.* (2009) 139:267–84. doi: 10.1016/j.cell.2009.09.028
100. National Research Council; Institute for Laboratory Animal Research, Commission on Life Sciences, Committee on Pain and Distress in Laboratory Animals. *Recognition and Alleviation of Pain and Distress in Laboratory Animals.* Washington, DC: National Academies Press (US) (1992). doi: 10.17226/1542
101. Miller RE, Malfait A-M. Osteoarthritis pain: what are we learning from animal models? *Best Pract Res Clin Rheumatology.* (2017) 31:676–87. doi: 10.1016/j.berh.2018.03.003
102. Fillingim RB, Loeser JD, Baron R, Edwards RR. Assessment of chronic pain: domains, methods, and mechanisms. *J Pain.* (2016) 17:T10–20. doi: 10.1016/j.jpain.2015.08.010
103. Woolf CJ. Central sensitization: implications for the diagnosis and treatment of pain. *Pain.* (2011) 152:S2–15. doi: 10.1016/j.jpain.2010.09.030
104. Bazzari AH, Bazzari FH. Advances in targeting central sensitization and brain plasticity in chronic pain. *Egypt J Neurol Psychiatry Neurosurg.* (2022) 58:38. doi: 10.1186/s41983-022-00472-y
105. Scher C, Meador L, Cleave JHV, Reid MC. Moving beyond pain as the fifth vital sign and patient satisfaction scores to improve pain care in the 21st century. *Pain Manag Nurs.* (2018) 19:125–9. doi: 10.1016/j.pmn.2017.10.010
106. Naye F, Cachinho C, Tremblay A-P, Lavoie MS-G, Lepage G, Larochelle E, et al. How to objectively assess and observe maladaptive pain behaviors in clinical rehabilitation: a systematic search and review. *Arch Physiother.* (2021) 11:15. doi: 10.1186/s40945-021-00109-y
107. Sharp Y, Tabor G. An investigation into the effects of changing dorso-plantar hoof balance on equine hind limb posture. *Animals Open Access J MDPI.* (2022) 12:3275. doi: 10.3390/ani12233275
108. Mawldsey A, Kelly EP, Smith FH, Brophy PO. Linear assessment of the thoroughbred horse: an approach to conformation evaluation. *Equine Vet J.* (1996) 28:461–7. doi: 10.1111/j.2042-3306.1996.tb01618.x
109. Carini F, Mazzola M, Fici C, Palmeri S, Messina M, Damiani P, et al. Posture and posturology, anatomical and physiological profiles: overview and current state of art. *Acta Biomed.* (2017) 88:11–6. doi: 10.23750/abm.v88i1.5309
110. Anker LC, Weerdesteijn V, Nes van IJW, Nienhuis B, Straatman H, Geurts ACH. The relation between postural stability and weight distribution in healthy subjects. *Gait Posture.* (2008) 27:471–7. doi: 10.1016/j.gaitpost.2007.06.002
111. Buchner HHF, Obermüller S, Scheidl M. Body centre of mass movement in the sound horse. *Vet J.* (2000) 160:225–34. doi: 10.1053/tvj.2000.0507
112. Ross MW. Lameness in horses: basic facts before starting. In: Ross MW, Dyson SJ, editors. *Diagnosis and management of lameness in the horse.* St Louis: Elsevier (2010). p. 3–8.
113. Buchner HH, Obermüller S, Scheidl M. Body centre of mass movement in the lame horse. *Equine Vet J Suppl.* (2001) 33:122–7. doi: 10.1111/j.2042-3306.2001.tb05374.x
114. Lesimple C, Fureix C, Aube L, Hausberger M. Detecting and measuring back disorders in nonverbal individuals: the example of domestic horses. *Animal Behav Cogn.* (2016) 3:159–79. doi: 10.12966/abc.05.08.2016
115. Fureix C, Hausberger M, Seneque E, Morisset S, Baylac M, Cornette R, et al. Geometric morphometrics as a tool for improving the comparative study of behavioural postures. *Naturwissenschaften.* (2011) 98:583. doi: 10.1007/s00114-011-0803-2