

RESEARCH

Open Access



# Comparison of a multiplex PCR with DNA barcoding for identification of container breeding mosquito species

Julia Reichl<sup>1,2</sup>, Christina Prosegger<sup>2</sup>, Sarah Petutschnig<sup>1</sup>, Maria Sophia Unterköfler<sup>2</sup>, Karin Bakran-Lebl<sup>1</sup>, Mateusz Markowicz<sup>1</sup>, Alexander Indra<sup>1</sup> and Hans-Peter Fuehrer<sup>2\*</sup>

## Abstract

**Background** Identification of mosquitoes greatly relies on morphological specification. Since some species cannot be distinguished reliably by morphological methods, it is important to incorporate molecular techniques into the diagnostic pipeline. DNA barcoding using Sanger sequencing is currently widely used for identification of mosquito species. However, this method does not allow detection of multiple species in one sample, which would be important when analysing mosquito eggs. Detection of container breeding *Aedes* is typically performed by collecting eggs using ovitraps. These traps consist of a black container filled with water and a wooden spatula inserted for oviposition support. *Aedes* mosquitoes of different species might lay single or multiple eggs on the spatula. In contrast to Sanger sequencing of specific polymerase chain reaction (PCR) products, multiplex PCR protocols targeting specific species of interest can be of advantage for detection of multiple species in the same sample.

**Methods** For this purpose, we adapted a previously published PCR protocol for simultaneous detection of four different *Aedes* species that are relevant for Austrian monitoring programmes, as they can be found in ovitraps: *Aedes albopictus*, *Aedes japonicus*, *Aedes koreicus*, and *Aedes geniculatus*. For evaluation of the multiplex PCR protocol, we analysed 2271 ovitrap mosquito samples from the years 2021 and 2022, which were collected within the scope of an Austrian nationwide monitoring programme. We compared the results of the multiplex PCR to the results of DNA barcoding.

**Results** Of 2271 samples, the multiplex PCR could identify 1990 samples, while species determination using DNA barcoding of the mitochondrial cytochrome c oxidase subunit I gene was possible in 1722 samples. The multiplex PCR showed a mixture of different species in 47 samples, which could not be detected with DNA barcoding.

**Conclusions** In conclusion, identification of *Aedes* species in ovitrap samples was more successful when using the multiplex PCR protocol as opposed to the DNA barcoding protocol. Additionally, the multiplex PCR allowed us to detect multiple species in the same sample, while those species might have been missed when using DNA barcoding with Sanger sequencing alone. Therefore, we propose that the multiplex PCR protocol is highly suitable and of great advantage when analysing mosquito eggs from ovitraps.

**Keywords** Multiplex PCR, *Aedes*, *Aedes albopictus*, Mosquitoes, Austria, Monitoring, Identification

\*Correspondence:  
Hans-Peter Fuehrer  
hans-peter.fuehrer@vetmeduni.ac.at  
Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Background

Identification of mosquito species has become increasingly important in the last decades, due to the spread of (potentially) invasive species [1]. Container breeding mosquito species, which use natural or artificial containers for oviposition [2], are of particular concern due to their potential to be competent vectors for a variety of pathogens. The increase in global transport and traffic enables the introduction and spread of those container breeding mosquitoes worldwide while climate change and concomitant warmer temperatures facilitate establishment of stable populations [3, 4]. Additionally, container breeding *Aedes* mosquitoes can lay eggs that can withstand desiccation and cold temperatures, which allows overwintering in temperate regions, such as Europe [5–7]. In the last decades, three container breeding *Aedes* species have become particularly important for Europe, namely the Asian bush mosquito (*Aedes japonicus*), the Korean bush mosquito (*Aedes koreicus*) and the Asian tiger mosquito (*Aedes albopictus*).

*Aedes albopictus*, the Asian tiger mosquito, is the most relevant mosquito of these three for human and veterinary health. It is a competent vector for viruses, such as dengue, Zika and chikungunya [8–10], as well as for filarial helminths, namely *Dirofilaria repens* and *Dirofilaria immitis* [11–14]. The native regions of *Ae. albopictus* are the tropical and subtropical Asian-Pacific regions from where it spread over all continents except the Antarctica [6, 8]. In Europe, the Asian tiger mosquito was first found in Albania in 1979 and later in Italy in 1990/1991. Since then, it has spread to more than 25 European countries [15–18].

*Aedes japonicus* originates from temperate regions of Eastern Asia [19] and was first reported in Europe in 2000 in northern France [20]. Under laboratory conditions, it is able to transmit viruses such as Japanese encephalitis virus [21, 22] and La Crosse virus [23], while West Nile virus has been identified multiple times in field-collected *Ae. japonicus* mosquitoes in the USA [24, 25]. Additionally, Austria reported the first natural infection of field-collected *Ae. japonicus* with Usutu virus in 2019 [26].

The Korean bush mosquito is native to Korea, Japan, north-eastern China and the far eastern parts of Russia [19, 27–29]. In 2008, *Ae. koreicus* was first found in Europe, in eastern Belgium [30]. Since then, a few other European countries, such as Italy [31, 32], Germany [33, 34], Switzerland [35], Hungary [36], the European part of Russia [37] and Austria [38] reported the presence of *Ae. koreicus*. The role of *Ae. koreicus* in transmission of pathogens remains unclear. However, studies suggest that this species is able to transmit Japanese encephalitis and *Dirofilaria immitis*, as well as some viruses such as chikungunya, under laboratory conditions [39–42].

In Austria, the first report of an exotic *Aedes* species showed the presence of *Ae. japonicus* in Styria in 2011 [43]. *Aedes japonicus* has been found in all provinces in Austria since its first report, has established stable populations, and can now be found in high numbers in all parts of Austria [44, 45]. *Aedes albopictus* was the second exotic *Aedes* species to be reported in Austria. It was first identified in Burgenland in 2012 [43]. After some initial single reports of *Ae. albopictus* mainly in Western Austria along the motorways, the number of reports has rapidly increased recently [38, 45]. In 2020, *Ae. albopictus* was first identified in Vienna, the capital city of Austria [46], and was found in the city of Graz in 2021 [47]. In 2022, the Asian tiger mosquito was found in every province in Austria if highway stations are included. However, stable populations are currently only documented in Vienna and Graz [48]. *Aedes koreicus* was first identified in Austria in 2017 in Carinthia [49]. Since then, there were some single reports in Carinthia, Styria and Tyrol [38, 48].

Identification of mosquitoes and their eggs greatly relies on morphological examination under a stereo microscope. Depending on the stage, some species might not be distinguishable with morphological methods. Therefore, it is important to incorporate molecular techniques for identification of mosquitoes [50–52]. DNA barcoding uses polymerase chain reaction (PCR) techniques to amplify conserved regions of the DNA with subsequent sequencing of the amplicon. The sequence obtained can then be compared with reference sequences in databases such as NCBI GenBank [53] to identify the mosquito species [54]. The most common DNA barcode is the mitochondrial cytochrome c oxidase subunit I (mtCOI) gene, which can serve as a molecular marker as it generally shows a low level of intra-species variation, but a high level of inter-species variation [54, 55]. While DNA barcoding using Sanger sequencing can additionally help with identifying cryptic species [56, 57], it does not allow accurate identification of multiple species in one sample [58]. Collection of *Aedes* eggs from container breeding species is typically performed by using so-called ovitraps, which consist of a black container filled with water and a wooden spatula plunged inside for oviposition support [44]. *Aedes* mosquitoes of different species might lay single or multiple eggs on the spatula. Morphological differentiation is possible but might be difficult depending on the species [59]. As molecular identification with Sanger sequencing of the mtCOI gene does not allow for identification of multiple species in one sample [58], multiplex PCR protocols targeting the specific species of interest can be of advantage for this purpose.

Bang et al. [60] designed a multiplex PCR protocol for simultaneous detection of six different *Aedes* species: *Ae.*

*albopictus*, *Ae. koreicus*, *Ae. japonicus*, *Ae. flavopictus*, *Ae. togoi* and *Ae. hatorii*. As only the first three species are relevant for Austria, we adapted the PCR protocol and further developed it so that identification of an Austrian native *Aedes* species, *Ae. geniculatus*, was also possible. Ovitrap mosquito samples from the years 2021 and 2022 deriving from an Austrian nationwide monitoring program [48, 61] were used for evaluation of the adapted multiplex PCR protocol and a comparison with morphological identification and DNA barcoding of the mtCOI gene.

Methods

Mosquito sampling

*Aedes* eggs were collected over the course of 2 years (2021, 2022) within the scope of a nationwide mosquito monitoring program. Ovitrap, representing the ideal habitat for breeding of *Aedes* mosquitoes, were set up in all nine provinces of Austria from the beginning of May until the end of October in both years. Black plastic containers (1 L) filled with tap water (approximately 0.75 L) served as ovitraps. For oviposition possibility, a wooden spatula, fixed with a stainless-steel clamp, was inserted into the water. The wooden spatula and the water were exchanged on a weekly basis, and the removed spatula was sent to the laboratory for further analysis.

Morphological analysis

Wooden spatulas were examined under the stereo microscope for the presence of mosquito eggs, more specifically *Aedes* eggs. Eggs were identified morphologically to species level, if possible, and subsequently removed from the spatula and collected in an Eppendorf tube (1.5 mL). Tubes were stored at -80 °C until molecular analysis was performed.

DNA extraction

For molecular analysis, all eggs from each wooden spatula were homogenized using one ceramic bead (2.8 mm Precellys Ceramic Beads, VWR, Darmstadt, Germany) and a TissueLyser II (Qiagen, Hilden, Germany) as described previously [38]. DNA of the samples from 2021 was extracted by the University of Veterinary Medicine, Vienna using the innuPREP DNA Mini Kit (Analytik Jena, Jena, Germany) according to the manufacturer's instructions. In 2022, molecular analyses were performed by the Austrian Agency for Health and Food Safety, which is why the protocol was changed and adapted to the available equipment at that laboratory. DNA isolation of the samples from 2022 was carried out with the BioExtract SuperBall Kit (Biosellal, Dardilly, France) on a King-Fisher Flex96 robot (Thermo Fisher Scientific, Waltham, USA) following an in-house protocol.

Multiplex PCR

The universal forward primer (Aedes-F), as well as the specific reverse primers for *Ae. albopictus* (ALB-R), *Ae. japonicus* (JAP-R) and *Ae. koreicus* (KOR-R), targeting the ribosomal ITS2 region were taken from the original protocol [60]. For *Ae. geniculatus*, we used all ITS2 reference sequences from the NCBI GenBank database to design a reverse primer (GEN-R) specific to that species and compatible with the other primers of the multiplex PCR. Primer design was performed with the software CLC Genomics Workbench 10 (Qiagen, Hilden, Germany). Primer sequences and details are shown in Table 1.

PCR reactions consisted of 10 µL REDTaq Ready Mix (Merck, Darmstadt, Germany), 7 µL nuclease-free water, 0.4 µL of each primer (10 µM) and 1 µL of DNA, while the PCR conditions described by Bang et al. [60] were applied (94 °C/5 min; then 94 °C/30 s, 56 °C/30 s, 72 °C/30 s for 35 cycles; and the final extension at 72 °C for 5 min). Gel electrophoresis (gel: 2.0% agarose gel in 0.5× TBE buffer; voltage: 100 V; run time: 90–120 min) and visualisation with the Gel Doc 2000 (Bio-Rad Laboratories, Hercules, USA) was performed to identify potential PCR products. The GeneRuler 100 bp DNA Ladder (Thermo Fisher Scientific, Waltham, USA) was used for sizing of PCR products. For confirming correct PCR conditions and successful primer binding, the PCR was first evaluated with representative DNA samples of the four relevant *Aedes* species (identified by DNA barcoding and morphological analysis) before the protocol was applied to the collected samples. The representative samples included *Ae. koreicus* larvae from the Viennese Central Cemetery [62], while the egg samples of the other species were obtained during the nationwide monitoring program in Austria [48, 61].

DNA barcoding

DNA barcoding of the mitochondrial cytochrome oxidase subunit I (mtCOI) gene was performed as previously described [56]. In 2021, the PCR products amplified with the primers LepF1 (5'-ATTCAACCAATCATAAAG ATAT-3') and LepR1 (5'-TAAACTTCTGGATGTCCA

Table 1 Primers used for the multiplex PCR protocol

Primer	Sequence (5'-3')	Forward/reverse	Source
Aedes-F	AGGACACATGAACCCACA	Forward	[60]
ALB-R	GGAGCACTGAGAGTTCCA	Reverse	[60]
JAP-R	TATACTACGCTGCCGAGAGG	Reverse	[60]
KOR-R	GCCTACTGATTGACGGGGTA	Reverse	[60]
GEN-R	ATGTATTCACCAACCGGG	Reverse	This study

AAAA-3') were sequenced at LGC Genomics GmbH, Berlin, Germany.

In 2022, PCR amplification was performed at the Austrian Agency for Health and Food Safety with the primers M13F-LepF1 (5'-TGTAACGACGGCCAGATTC AACCAATCATAAAGATAT-3') and M13R-LepR1 (5'-CAGGAAACAGCTATGACTAACTTCTGGATGTC CAAAAA-3'). M13 sequences were added to the primers (without resulting in different primer binding properties) for facilitating the subsequent workflow. The PCR conditions of the protocol using the primers LepF1 and LepR1 were applied [56]. For PCR cleanup, 2 µL of the ExoSAP-IT Express reagent (Applied Biosystems, Waltham, USA) were mixed with 5 µL PCR product and incubated at 37 °C for 4 min and 80 °C for 1 min. Subsequent cycle sequencing reactions with the sequencing primer M13R (5'-CAGGAAACAGCTATGAC-3') were set up using the BigDye Terminator v3.1 Cycle Sequencing kit (Applied Biosystems, Waltham, USA). The reactions contained 8 µL BigDye Terminator 3.1 Ready Reaction Mix, 9.68 µL ddH<sub>2</sub>O, 0.32 µL of the sequencing primer M13R (10 µM) and 2 µL of template. The conditions for the cycle sequencing were chosen as recommended by the manufacturer. After cycle sequencing, the reactions were purified by adding 90 µL of SAM solution (Applied Biosystems, Waltham, USA) and 20 µL of the XTerminator solution (Applied Biosystems, Waltham, USA). After shaking for 20 min at 1800 rpm on an Eppendorf

MixMate (Eppendorf, Hamburg, Germany) and centrifugation at 1000g for 2 min, the sequencing reactions were analysed with an ABI Genetic Analyzer 3500 (PA Protocol: BDTv3.1\_PA\_Protocol\_POP7, Basecall Version: KB 1.4.1.8) (Applied Biosystems, Waltham, USA). After analysis, the resulting sequences were compared to sequences of two public databases (NCBI Genbank, [www.ncbi.nlm.nih.gov/genbank](http://www.ncbi.nlm.nih.gov/genbank); BOLD Systems, [www.boldsystems.org](http://www.boldsystems.org)) for species identification.


### Data analysis

All data obtained was documented in Microsoft Excel (Microsoft, Washington, USA) and analysed using the software R version 4.2.3 [63].

## Results

### Confirmation of PCR conditions and primer binding

For a first evaluation, a total of 31 DNA samples from ovitraps and larvae (8 *Ae. koreicus* larval samples, eight *Ae. albopictus* egg samples, eight *Ae. japonicus* egg samples, seven *Ae. geniculatus* egg samples) and one negative control (water) were analysed using the adapted multiplex PCR protocol. The samples were previously identified morphologically and by DNA barcoding. After gel electrophoresis (Fig. 1), all samples showed specific PCR products of the correct sizes according to the expected species (438 bp for *Ae. albopictus*, 361 bp for *Ae. koreicus*, 310 bp for *Ae. geniculatus* and 160 bp for *Ae. japonicus*).



**Fig. 1** Results of the multiplex PCR of representative DNA samples of the four relevant *Aedes* species. The 31 DNA samples (lanes 1–8: *Ae. albopictus* eggs; lanes 9–16: *Ae. japonicus* eggs; Lanes 17–24: *Ae. koreicus* larvae; lanes 25–31: *Ae. geniculatus* eggs) were previously identified morphologically and with DNA barcoding and subsequently analysed using the multiplex PCR protocol. All samples show specific PCR products of the correct sizes according to the expected species: 438 bp for *Ae. albopictus*, 361 bp for *Ae. koreicus*, 310 bp for *Ae. geniculatus* and 160 bp for *Ae. japonicus*. The sample in lane 3 (expected to be *Ae. albopictus*) shows an additional PCR product at 160 bp, indicating the presence of *Ae. japonicus*



One sample, which was expected to be *Ae. albopictus*, showed an additional PCR product at 160 bp, indicating the presence of *Ae. japonicus*.

### Analysis of ovitrap samples from 2021 and 2022

In total, 2271 samples from the years 2021 and 2022 were analysed using morphology, the adapted multiplex PCR and DNA barcoding with Sanger sequencing. Morphological specification of the eggs revealed *Ae. albopictus* in 203 samples and *Ae. geniculatus* in 122 samples, while 104 samples were either classified as another species or not identified at all. Differentiation of *Ae. japonicus* and *Ae. koreicus* was not possible when morphologically looking at egg samples, which is why 1750 samples were classified as *Ae. koreicus/Ae. japonicus*. In an additional 92 samples, morphological examination revealed the presence of more than one species (5 *Ae. albopictus/Ae. geniculatus*, 8 *Ae. albopictus/Ae. japonicus/Ae. koreicus*, 79 *Ae. geniculatus/Ae. japonicus/Ae. koreicus*).

The results of the multiplex PCR analyses identified 162 *Ae. albopictus*, 1705 *Ae. japonicus* and 76 *Ae. geniculatus* samples. In those samples, DNA of only one species was present. 281 samples could not be identified, as no PCR product was visible on the gel. *Aedes koreicus* was not detected in any of the analysed samples. According to the multiplex PCR, DNA of multiple species was present in 47 samples. Of those, 18 were mixtures of *Ae. albopictus* and *Ae. japonicus*, while 29 samples contained DNA of *Ae. japonicus* and *Ae. geniculatus*.

DNA barcoding of mtCOI detected 163 *Ae. albopictus*, 88 *Ae. geniculatus* and 1281 *Ae. japonicus*, while *Ae. koreicus* was not identified. A total of 190 samples were classified as another species or organism. In 549 samples, DNA barcoding did not show any results due to low sequencing quality or lack of amplification of mtCOI. With DNA barcoding using Sanger sequencing, mixtures of different species could not be detected (Table 2).

To compare the identification of container-breeding mosquitoes with DNA barcoding to the multiplex PCR protocol, contingency tables for the species *Ae. albopictus*, *Ae. japonicus* and *Ae. geniculatus* were created (Table 2). Due to the absence of *Ae. koreicus* in the analysed ovitrap samples, it was not possible to compare the two methods for this species. Percent agreement and Cohen's Kappa coefficients ( $\kappa$ ) were calculated to demonstrate the agreement between the two applied methods for each parameter/species: *Ae. albopictus* 97.58%,  $\kappa=0.83$  [95% confidence interval (CI) 0.78–0.87]; *Ae. japonicus* 76.62%,  $\kappa=0.50$  (95% CI 0.46–0.53); *Ae. geniculatus* 97.49%,  $\kappa=0.69$  (95% CI 0.62–0.77).

Samples that, according to the multiplex PCR, contained DNA of *Ae. albopictus* and *Ae. japonicus* ( $n=18$ ), were identified as *Ae. albopictus* in 10 cases and as *Ae.*

**Table 2** Contingency tables for the three species *Ae. albopictus*, *Ae. japonicus* and *Ae. geniculatus*

DNA barcoding		Multiplex PCR		
		Positive	Negative	Total
<i>Aedes albopictus</i>	Positive	144	19	163
	Negative	36	2072	2108
	Total	180	2091	2271
<i>Aedes japonicus</i>	Positive	1251	30	1281
	Negative	501	489	990
	Total	1752	519	2271
<i>Aedes geniculatus</i>	Positive	68	20	88
	Negative	37	2146	2183
	Total	105	2166	2271

The table shows the amounts of samples positive or negative for the individual species, analysed with multiplex PCR and DNA barcoding

*japonicus* in two cases when using DNA barcoding. Six of those samples showed no result with Sanger sequencing. While the multiplex PCR detected *Ae. japonicus* and *Ae. geniculatus* ( $n=29$ ), DNA barcoding showed no result ( $n=4$ ), *Ae. japonicus* ( $n=8$ ), *Ae. geniculatus* ( $n=15$ ) or another organism ( $n=2$ ).

The comparison of the molecular techniques with morphology was out of the scope of this study. However, to be able to better evaluate the performance of the two methods, additional contingency tables including morphological results were created for each species (Additional file 1: Tables S1, S2).

### Discussion

DNA barcoding using Sanger sequencing offers the possibility to identify a wide variety of different mosquito species (and other organisms). However, in contrast to a multiplex PCR, the method is not suitable for detection of multiple species in one sample. In a total of 47 samples, Sanger sequencing either detected only one species or did not show any result, while the multiplex PCR revealed the presence of multiple species. In 8 of 18 cases, *Ae. albopictus* was not identified in a sample containing more than one species when using DNA barcoding. Based on the results of this study, it can be assumed that using DNA barcoding with Sanger sequencing can lead to an under-reporting when dealing with samples that might contain DNA of several species. Hence, we propose that the described multiplex PCR protocol is of great advantage for certain monitoring purposes (e.g. ovitrap monitoring).

A Cohen's kappa coefficient between 0.40 and 0.59 indicates weak agreement, while values from 0.60–0.79

to 0.80–0.90 are to be interpreted as moderate and strong agreement, respectively [64]. Based on the Cohen's kappa coefficients, the agreement between the two methods for detecting *Ae. albopictus* is quite high, which is also shown by the percent agreement (97.58%). In contrast, Cohen's kappa coefficient of *Ae. japonicus* ( $\kappa=0.50$ ) indicates only little agreement between the PCR and DNA barcoding, demonstrated also by a lower percent agreement of 76.62%. The results further indicate moderate agreement between the two methods when looking at the identification of *Ae. geniculatus*, with a Cohen's kappa coefficient of 0.69 and a percent agreement of 97.49%. Differences in the degree of agreement between the species might be caused due to possible biases (primer bias, PCR selection, PCR drift), which result in higher amplification of some templates due to their properties (e.g. GC content and gene copy numbers) [65]. Another reason might be the difference in sample size, as a considerably higher number of samples positive for *Ae. japonicus* was analysed. Additionally, *Ae. japonicus* is considered a species complex, which is (currently) composed of four subspecies [66]. PCR and/or sequencing primers could show different binding properties related to the different subspecies, affecting the overall performance of those methods. Recently, it has been suggested that *Ae. geniculatus* either has cryptic sibling species or that it is also a species complex comprised of multiple subspecies [67]. However, this still needs to be further investigated since currently there are only few DNA sequences available in public databases for this species. This also posed a challenge for designing the primer for *Ae. geniculatus* in this study.

When comparing the number of negative results (no species identification), the multiplex PCR performed better than DNA barcoding. A total of 281 samples could not be identified by PCR, while DNA barcoding with Sanger sequencing did not lead to a result in 549 cases. Inspection of the contingency table (Table 2) makes it clear that the multiplex PCR generally performed slightly better in identifying the relevant container-breeding *Aedes* species. The biggest difference in performance can be observed with *Ae. japonicus*. A total of 501 PCR positive samples could not be identified as *Ae. japonicus* by Sanger sequencing, while only 30 *Ae. japonicus* samples would have been overlooked when using the multiplex PCR. Other organisms (e.g. Ceratopogonidae eggs or Amoebozoa such as *Vanella simplex*), which can also be detected during DNA barcoding, might be present in some of the samples and thereby lower the detection rate of sequencing when looking at the four relevant *Aedes* species.

While the aim of this study was primarily to compare the two molecular methods with each other, morphological results were included for a better overview

and evaluation of the techniques. Generally, morphological analysis resulted in less non-identifiable samples compared to the multiplex PCR and DNA barcoding. A reason for the weaker performance of the molecular techniques might be that there is not enough DNA present (e.g. in samples with few eggs or after errors during DNA extraction). Nonetheless, some samples would have been overlooked when using morphological inspection alone (especially if scientists are not experienced in morphological specification of mosquito eggs). Therefore, we suggest complementing morphological assessment with molecular techniques to decrease mis- or non-identifications. Due to its better performance and higher agreement with morphology, we propose to use the multiplex PCR as a second method for identification of eggs from ovitrap samples. The analysed ovitrap samples from 2021 to 2022 did not include any *Ae. koreicus* eggs. While larvae of *Ae. koreicus* were found in the Viennese Central Cemetery in 2021, ovitraps in the same region were negative for this species. Possible reasons for the absence of eggs might be that there were too few *Ae. koreicus* females present in the area or that they preferred other breeding sites [62]. A comparison between Sanger sequencing and the multiplex PCR protocol was therefore not possible for this species and needs to be investigated in further experiments. However, the proposed PCR protocol was validated with representative *Ae. koreicus* larval samples. As species-specific bands were observed on the agarose gel, it is highly likely that the PCR protocol is also suitable for identification of *Ae. koreicus* in ovitrap samples.

Differentiation of *Ae. koreicus* and *Ae. geniculatus* might be difficult in some cases, as the PCR products lie close together when being visualised by gel electrophoresis. It is therefore important to ensure that the run time of the gel electrophoresis is adequate (in our cases 1.5–2 h). Additionally, results of other methods (e.g. morphology) can be taken into account for evaluation of the PCR results. If interpretation of the results concerning *Ae. koreicus* and *Ae. geniculatus* is still not possible, a second PCR using the universal forward primer and only one reverse primer (specific to *Ae. koreicus* or *Ae. geniculatus*) can be carried out for confirmation. Alternatively, agarose gel electrophoresis can be substituted by a fragment analysis using a suitable instrument (e.g. ABI 3500 Genetic Analyzer, Applied Biosystems, Waltham, USA) and fluorescently labelled primers. Fragment analysis allows for separation of PCR products that show a size difference of only 1–2 nt and can thereby improve analysis and interpretation of multiplex PCR reactions. Furthermore, the protocol can be adapted for analysis with a real-time PCR (RT-PCR) machine using fluorescently labelled primers or probes. Additionally, a RT-PCR approach offers higher

sensitivity for detecting the four species and provides results more quickly than a conventional PCR as there is no need for subsequent gel electrophoresis. Additionally, PCR products can be quantified when using RT-PCR [68]. In the future, mosquito species identification will most probably benefit from the recent emergence of next-generation sequencing applications. Using NGS technology, DNA metabarcoding can be performed, which allows detection of multiple species in the same sample while being able to identify a wide range of different species. At the moment, however, PCR approaches still score with less time-consuming protocols and fewer costs, when compared with NGS applications. Nonetheless, further development of new methods for mosquito identification (and pathogen detection) would likely improve current workflows.

## Conclusions

The multiplex PCR protocol described in this study can serve as a powerful tool for large-scale analysis of container-breeding mosquitoes. Especially when detection of multiple species in the same sample might be necessary (e.g. ovitrap samples), the multiplex PCR is more suitable than DNA barcoding with Sanger sequencing. Additionally, PCR analyses are typically less time-consuming and cheaper compared to sequencing technologies. Therefore, we propose that this multiplex PCR protocol is highly suitable and of great advantage for analysing container-breeding mosquitoes.

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13071-024-06255-z>.

**Additional file 1: Table S1.** Contingency table for the three species *Ae. albopictus*, *Ae. japonicus* and *Ae. geniculatus* for comparison of the multiplex PCR and morphological analysis. The samples positive or negative for the individual species, when examined morphologically and by the multiplex PCR, are depicted in this table. **Table S2.** Contingency table for the three species *Ae. albopictus*, *Ae. japonicus* and *Ae. geniculatus* for comparison of DNA barcoding and morphological analysis. The amount of positive and negative samples after DNA barcoding and morphological examination are shown in this table.

## Acknowledgements

We wish to thank all citizen scientists and employees from AGES who conducted the sampling of the ovitraps.

## Author contributions

HPF, KBL and AI supervised the study. JR, CP, SP and MSU performed molecular genetic laboratory work. KBL coordinated the sampling and carried out morphological analysis. MM acquired funding via the ESMOS project. JR wrote the first draft. All authors have reviewed the final manuscript.

## Funding

Financial support was provided by the Climate and Energy Fund of the Austrian Federal Government via the ESMOS project (FFG 901442) within the framework of the Austrian Climate Research Programme (ACRP), 15th call.

## Availability of data and materials

The datasets used and/or analysed during the current study are included in this published article (and its supplementary information files).

## Declarations

### Ethics approval and consent to participate

Not applicable.

### Consent for publication

Not applicable.

### Competing interests

The authors declare that they have no competing interests.

### Author details

<sup>1</sup>Institute for Medical Microbiology and Hygiene, Austrian Agency for Health and Food Safety (AGES), Vienna, Austria. <sup>2</sup>Institute of Parasitology, Department of Pathobiology, University of Veterinary Medicine Vienna, Vienna, Austria.

Received: 22 December 2023 Accepted: 20 March 2024

Published online: 02 April 2024

## References

- Hoque MM, Valentine MJ, Kelly PJ, Barua S, Murillo DFB, Wang C. Modification of the Folmer primers for the cytochrome c oxidase gene facilitates identification of mosquitoes. *Parasit Vectors*. 2022;15:437.
- Reinbold-Wasson DD, Reiskind MH. Comparative skip-oviposition behavior among container breeding *Aedes* spp. mosquitoes (Diptera: Culicidae). *J Med Entomol*. 2021;58:2091–100.
- Cunze S, Kochmann J, Koch LK, Klimpel S. *Aedes albopictus* and its environmental limits in Europe. *PLoS ONE*. 2016;11:e0162116.
- Eritja R, Palmer JRB, Roiz D, Sanpera-Calbet I, Bartumeus F. Direct evidence of adult *Aedes albopictus* dispersal by car. *Sci Rep*. 2017;7:14399.
- Juliano SA, Lounibos LP. Ecology of invasive mosquitoes: effects on resident species and on human health. *Ecol Lett*. 2005;8:558–74.
- Paupy C, Delatte H, Bagay L, Corbel V, Fontenille D. *Aedes albopictus*, an arbovirus vector: from the darkness to the light. *Microbes Infect*. 2009;11:1177–85.
- Tippelt L, Werner D, Kampen H. Tolerance of three *Aedes albopictus* strains (Diptera: Culicidae) from different geographical origins towards winter temperatures under field conditions in northern Germany. *PLoS ONE*. 2019;14:e0219553.
- Bonizzoni M, Gasperi G, Chen X, James AA. The invasive mosquito species *Aedes albopictus*: current knowledge and future perspectives. *Trends Parasitol*. 2013;29:460–8.
- Gloria-Soria A, Payne AF, Bialosuknia SM, Stout J, Mathias N, Eastwood G, et al. Vector competence of *Aedes albopictus* populations from the Northeastern United States for chikungunya, dengue, and Zika viruses. *Am J Trop Med Hyg*. 2020;104:1123–30.
- Medlock JM, Hansford KM, Schaffner F, Versteirt V, Hendrickx G, Zeller H, et al. A review of the invasive mosquitoes in Europe: ecology, public health risks, and control options. *Vector Borne Zoonotic Dis*. 2012;12:435–47.
- Cancrini G, Romi R, Gabrielli S, Toma L, M DIP, Scaramozzino P. First finding of *Dirofilaria repens* in a natural population of *Aedes albopictus*. *Med Vet Entomol*. 2003;17:448–51.
- Fuehrer HP, Auer H, Leschnik M, Silberman K, Duschner G, Joachim A. *Dirofilaria* in humans, dogs, and vectors in Austria (1978–2014)-from imported pathogens to the endemicity of *Dirofilaria repens*. *PLoS Negl Trop Dis*. 2016;10:e0004547.
- Fuehrer HP, Morelli S, Unterkofler MS, Bajer A, Bakran-Lebl K, Dwuznik-Szarek D, et al. *Dirofilaria* spp. and *Angiostrongylus vasorum*: current risk of spreading in Central and Northern Europe. *Pathogens*. 2021;10:1268 <https://doi.org/10.3390/pathogens10101268>.

14. Nayar JK, Knight JW. *Aedes albopictus* (Diptera: Culicidae): an experimental and natural host of *Dirofilaria immitis* (Filarioidea: Onchocercidae) in Florida, USA. *J Med Entomol*. 1999;36:441–8.
15. Adhami J, Reiter P. Introduction and establishment of *Aedes (Stegomyia) albopictus* skuse (Diptera: Culicidae) in Albania. *J Am Mosq Control Assoc*. 1998;14:340–3.
16. ECDC. *Aedes albopictus* - current known distribution: March 2022: European Centre for Disease Prevention and Control. 2022. <https://www.ecdc.europa.eu/en/publications-data/aedes-albopictus-current-known-distribution-march-2022>.
17. Medlock JM, Hansford KM, Versteirt V, Cull B, Kampen H, Fontenille D, et al. An entomological review of invasive mosquitoes in Europe. *Bull Entomol Res*. 2015;105:637–63.
18. Battaglia V, Agostini V, Moroni E, Colombo G, Lombardo G, Rambaldi Migliore N, et al. The worldwide spread of *Aedes albopictus*: new insights from mitogenomes. *Front Genet*. 2022;13:931163.
19. A revision of the adult and larval mosquitoes of Japan (including the Ryukyu Archipelago and the Ogasawara Islands) and Korea (Diptera: Culicidae). *Mosquito News*. 1980;40:2.
20. Schaffner F, Chouin S, Guilleateau J. First record of *Ochlerotatus (Finlaya) japonicus japonicus* (Theobald, 1901) in metropolitan France. *J Am Mosq Control Assoc*. 2003;19:1–5.
21. Faizah AN, Kobayashi D, Amoa-Bosompem M, Higa Y, Tsuda Y, Itokawa K, et al. Evaluating the competence of the primary vector, *Culex tritaeniorhynchus*, and the invasive mosquito species, *Aedes japonicus japonicus*, in transmitting three Japanese encephalitis virus genotypes. *PLoS Negl Trop Dis*. 2020;14:e0008986.
22. Van den Eynde C, Sohler C, Matthijs S, De Regge N. Japanese encephalitis virus interaction with mosquitoes: a review of vector competence, vector capacity and mosquito immunity. *Pathogens*. 2022;11:317. <https://doi.org/10.3390/pathogens11030317>.
23. Harris MC, Dotseth EJ, Jackson BT, Zink SD, Marek PE, Kramer LD, et al. La Crosse virus in *Aedes japonicus japonicus* mosquitoes in the Appalachian region, United States. *Emerg Infect Dis*. 2015;21:646–9.
24. DeCarlo CH, Campbell SR, Bigler LL, Mohammed HO. *Aedes japonicus* and West Nile virus in New York. *J Am Mosq Control Assoc*. 2020;36:261–3.
25. Centers for Disease Control and Prevention. Mosquito species in which West Nile virus has been detected, United States, 1999–2016. Centers for Disease Control and Prevention; 2016.
26. Camp JV, Kolodziejek J, Nowotny N. Targeted surveillance reveals native and invasive mosquito species infected with Usutu virus. *Parasit Vectors*. 2019;12:46.
27. Key to the Insects of the Russian Far East. Vol. IV Diptera and fleas. Part 2. Vladivostok: Dal'nauka; 2001.
28. Knight KL. Contributions to the mosquito fauna of Southeast Asia.—IV. Species of the subgroup chrysolineatus of group D, genus *Aedes*, subgenus *Finlaya* Theobald. *Contribut Am Entomol Inst*. 1968;2:1–45.
29. Ward RA. Fauna of the U.S.S.R. Diptera, Vol. III, No. 4, mosquitoes family culicidae. *Bullet Entomol Soc Am*. 1975;21:147.
30. Versteirt V, De Clercq EM, Fonseca DM, Pecor J, Schaffner F, Coosemans M, et al. Bionomics of the established exotic mosquito species *Aedes koreicus* in Belgium, Europe. *J Med Entomol*. 2012;49:1226–32.
31. Capelli G, Drago A, Martini S, Montarsi F, Soppelsa M, Delai N, et al. First report in Italy of the exotic mosquito species *Aedes (Finlaya) koreicus*, a potential vector of arboviruses and filariae. *Parasit Vectors*. 2011;4:188.
32. Negri A, Arnoldi I, Brilli M, Bandi C, Gabrieli P, Epis S. Evidence for the spread of the alien species *Aedes koreicus* in the Lombardy region, Italy. *Parasit Vectors*. 2021;14:534.
33. Werner D, Zielke DE, Kampen H. First record of *Aedes koreicus* (Diptera: Culicidae) in Germany. *Parasitol Res*. 2016;115:1331–4.
34. Pfützner WP, Lehner A, Hoffmann D, Czajka C, Becker N. First record and morphological characterization of an established population of *Aedes (Hulecoeteomyia) koreicus* (Diptera: Culicidae) in Germany. *Parasit Vectors*. 2018;11:662.
35. Suter T, Flacio E, Farina BF, Engeler L, Tonolla M, Muller P. First report of the invasive mosquito species *Aedes koreicus* in the Swiss-Italian border region. *Parasit Vectors*. 2015;8:402.
36. Kurucz K, Kiss V, Zana B, Schmieder V, Kepner A, Jakab F, et al. Emergence of *Aedes koreicus* (Diptera: Culicidae) in an urban area, Hungary, 2016. *Parasitol Res*. 2016;115:4687–9.
37. Ganushkina LA, Patraman IV, Rezza G, Migliorini L, Litvinov SK, Sergiev VP. Detection of *Aedes aegypti*, *Aedes albopictus*, and *Aedes koreicus* in the Area of Sochi, Russia. *Vector Borne Zoonotic Dis*. 2016;16:58–60.
38. Fuehrer HP, Schoener E, Weiler S, Barogh BS, Zittira C, Walder G. Monitoring of alien mosquitoes in Western Austria (Tyrol, Austria, 2018). *PLoS Negl Trop Dis*. 2020;14:e0008433.
39. Ganassi S, De Cristofaro A, Di Criscio D, Petrarca S, Leopardi C, Guarnieri A, et al. The new invasive mosquito species *Aedes koreicus* as vector-borne diseases in the European area, a focus on Italian region: what we know from the scientific literature. *Front Microbiol*. 2022;13:931994.
40. Montarsi F, Ciocchetta S, Devine G, Ravagnan S, Mutinelli F, di Frangipane Regalbono A, et al. Development of *Dirofilaria immitis* within the mosquito *Aedes (Finlaya) koreicus*, a new invasive species for Europe. *Parasit Vectors*. 2015;8:177.
41. Jansen S, Cadar D, Luhken R, Pfützner WP, Jost H, Oerther S, et al. Vector competence of the invasive mosquito species *Aedes koreicus* for arboviruses and interference with a novel insect specific virus. *Viruses*. 2021;12:2507. <https://doi.org/10.3390/v13122507>.
42. ECDC. *Aedes koreicus* - Factsheet for experts: European Centre for Disease Prevention and Control. 2014. <https://www.ecdc.europa.eu/en/disease-vectors/facts/mosquito-factsheets/aedes-koreicus>.
43. Seidel B, Duh D, Nowotny N, Allerberger F. Erstnachweis der Stechmücken *Aedes (Ochlerotatus) japonicus japonicus* (Theobald, 1901) in Österreich und Slowenien in 2011 und für *Aedes (Stegomyia) albopictus* (Skuse, 1895) in Österreich 2012 (Diptera: Culicidae). *Entomol Z*. 2012;122:223–6.
44. Bakran-Lebl K, Pree S, Brenner T, Daroglou E, Eigner B, Griesbacher A, et al. First nationwide monitoring program for the detection of potentially invasive mosquito species in Austria. *Insects*. 2022;13:276. <https://doi.org/10.3390/insects13030276>.
45. Schoener E, Zittira C, Weiss S, Walder G, Barogh BS, Weiler S, et al. Monitoring of alien mosquitoes of the genus *Aedes* (Diptera: Culicidae) in Austria. *Parasitol Res*. 2019;118:1633–8.
46. Bakran-Lebl K, Zittira C, Harl J, Shahi-Barogh B, Gratzl A, Ebmer D, et al. Arrival of the Asian tiger mosquito, *Aedes albopictus* (Skuse, 1895) in Vienna, Austria and initial monitoring activities. *Transbound Emerg Dis*. 2021;68:3145–50.
47. Reichl J, Prossegger C, Eichholzer B, Plauder P, Unterkofler MS, Bakran-Lebl K, et al. A citizen science report-tiger mosquitoes (*Aedes albopictus*) in allotment gardens in Graz, Styria, Austria. *Parasitol Res*. 2023;123:79.
48. Bakran-Lebl K, Reichl J. Ovitrap-Monitoring gebietsfremder Gelsenarten in Österreich - Jahresbericht 2022. Vienna: AGES - Austrian Agency for Health and Food Safety, Institute for Medical Microbiology und Hygiene DfVBD; 2023.
49. Seidel B, Hufnagl P, Nowotny N, Allerberger F, Indra A. Long-term field study on mosquitoes in Austria, in particular the invasive Korean bush-mosquito *Aedes (Finlaya) koreicus* (Edwards, 1917). *Beiträge zur Entomofaunistik*. 2020;21:237–40.
50. Gunay F, Alten B, Simsek F, Aldemir A, Linton YM. Barcoding Turkish *Culex* mosquitoes to facilitate arbovirus vector incrimination studies reveals hidden diversity and new potential vectors. *Acta Trop*. 2015;143:112–20.
51. Weeraratne TC, Surendran SN, Parakrama Karunaratne SHP. DNA barcoding of morphologically characterized mosquitoes belonging to the subfamily Culicinae from Sri Lanka. *Parasit Vectors*. 2018;11:266.
52. Hernandez-Triana LM, Brugman VA, Nikolova NI, Ignacio RA, Barrero E, Thorne L, et al. DNA barcoding of British mosquitoes (Diptera, Culicidae) to support species identification, discovery of cryptic genetic diversity and monitoring invasive species. *Zookeys*. 2019;832:57–76.
53. Sayers EW, Bolton EE, Brister JR, Canese K, Chan J, Comeau DC, et al. Database resources of the national center for biotechnology information. *Nucleic Acids Res*. 2022;50:D20–6.
54. Hebert PD, Cywinska A, Ball SL, deWaard JR. Biological identifications through DNA barcodes. *Proc Biol Sci*. 2003;270:313–21.
55. Hebert PD, Ratnasingham S, deWaard JR. Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. *Proc Biol Sci*. 2003;270:596–9.
56. Hebert PD, Penton EH, Burns JM, Janzen DH, Hallwachs W. Ten species in one: DNA barcoding reveals cryptic species in the neotropical skipper butterfly *Astraptes fulgerator*. *Proc Natl Acad Sci USA*. 2004;101:14812–7.
57. Ruiz-Lopez F, Wilkerson RC, Conn JE, McKeon SN, Levin DM, Quinones ML, et al. DNA barcoding reveals both known and novel taxa in the *Albitarsis*



- group (*Anopheles Nyssorhynchus*) of neotropical malaria vectors. *Parasit Vectors*. 2012;5:44.
58. Batovska J, Cogan NO, Lynch SE, Blacket MJ. Using next-generation sequencing for DNA barcoding: capturing allelic variation in ITS2. *G3*. 2017;7:19–29.
  59. Anicic N, Steigmiller K, Renaux C, Ravasi D, Tanadini M, Flacio E. Optical recognition of the eggs of four *Aedine* mosquito species (*Aedes albopictus*, *Aedes geniculatus*, *Aedes japonicus*, and *Aedes koreicus*). *PLoS ONE*. 2023;18:e0293568.
  60. Bang WJ, Won MH, Cho ST, Ryu J, Choi KS. A multiplex PCR assay for six *Aedini* species, including *Aedes albopictus*. *Parasit Vectors*. 2021;14:380.
  61. Bakran-Lebl K. Ovitrap-Monitoring gebietsfremder Gelsenarten in Österreich - Jahresbericht 2021. Vienna: AGES - Austrian Agency for Health and Food Safety, Institute for Medical Microbiology und Hygiene DfVBD; 2022.
  62. Unterkofler MS, Schreier L, Wiesbauer U, Zittra C, Grätzl A, Prosegger C, et al. Nachweis von heimischen und potenziell invasiven Stechmücken am Wiener Zentralfriedhof im Jahr 2021. *Entomologica Austriaca*. 2022;359–60.
  63. R Core Team. R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing; 2021.
  64. McHugh ML. Interrater reliability: the kappa statistic. *Biochem Med*. 2012;22:276–82.
  65. Acinas SG, Sarma-Rupavtarm R, Klepac-Ceraj V, Polz MF. PCR-induced sequence artifacts and bias: insights from comparison of two 16S rRNA clone libraries constructed from the same sample. *Appl Environ Microbiol*. 2005;71:8966–9.
  66. Kaufman MG, Fonseca DM. Invasion biology of *Aedes japonicus japonicus* (Diptera: Culicidae). *Annu Rev Entomol*. 2014;59:31–49.
  67. Prudhomme J, Fontaine A, Lacour G, Gantier JC, Diancourt L, Velo E, et al. The native European *Aedes geniculatus* mosquito species can transmit chikungunya virus. *Emerg Microbes Infect*. 2019;8:962–72.
  68. Kralik P, Ricchi M. A basic guide to real time PCR in microbial diagnostics: definitions, parameters, and everything. *Front Microbiol*. 2017;8:108.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.