

CASE REPORT

Open Access



Isolation and characterization of *Leptospira licerasiae* in Austrian swine — a first-time case report in Europe

Adi Steinrigl^{1*}, Denise Willixhofer¹, Martin Schindler¹, Susanne Richter¹, Christine Unterweger², Ahmed A. Ahmed³, Hans van der Linden³, Daniel R. Mende⁴, Nicholas Pucci⁴ and Romana Steinparzer¹

Abstract

Background Leptospiraceae comprise a diverse family of spirochetal bacteria, of which many are involved in infectious diseases of animals and humans. Local leptospiral diversity in domestic animals is often poorly understood. Here we describe the incidental detection of *Leptospira* (*L.*) *licerasiae* in an Austrian pig.

Case presentation During an experiment to characterize the pathogenesis of *L. interrogans* serovar Icterohaemorrhagiae in pigs, cultivation of a urine sample from a non-challenged contact pig resulted in growth of a spirochetal bacterium that tested negative for pathogenic *Leptospira* (LipL32 gene). PCR, Sanger sequencing and standard serotyping further confirmed that the recovered isolate was clearly different from the challenge strain *L. interrogans* serovar Icterohaemorrhagiae used in the animal experiment. Whole genome sequencing revealed that the isolate belongs to the species *L. licerasiae*, a tropical member of the Leptospiraceae, with no prior record of detection in Europe.

Conclusions This is the first report describing the occurrence of *L. licerasiae* in Europe. Since *L. licerasiae* is considered to have intermediate pathogenicity, it will be important to follow the geographical distribution of this species and its pathogenic and zoonotic potential in more detail.

Keyword *Leptospira licerasiae*, Pig, Austria, Whole genome sequencing

Background

The genus *Leptospira* currently comprises at least 68 different species, based on their whole genome sequence [1, 2]. These species can be further differentiated according to their pathogenic phenotype into pathogenic, intermediate, and saprophytic species [3, 4]. Thorough bioinformatics analysis enabled further distinction into four clades (P1, P2, S1 and S2) [2]. Besides genetic nomenclature, *Leptospira* are classified according to agglutination with reference antisera in the microscopic agglutination test (MAT) [5]. Thus, 300 different serovars can be distinguished which do, however, not align with the genetic classification system [5, 6]. Pathogenic *Leptospira* play an important role as causative disease agents in man and animals alike and leptospirosis has been acknowledged as the most widespread zoonosis in the world [7]. The

*Correspondence:

Adi Steinrigl
adi.steinrigl@ages.at

¹ Austrian Agency for Health and Food Safety, Institute for Veterinary Disease Control Moedling, Robert Koch Gasse 17, Mödling 2340, Austria

² Clinical Centre for Farm Animals and Food System Science, University of Veterinary Medicine, Veterinärplatz 1, Vienna 1221, Austria

³ Expertise Centre for Reference and Research On Leptospirosis/WOAH Reference Laboratory for Leptospirosis, Department of Medical Microbiology and Infection Prevention, Amsterdam University Medical Centers, Meibergdreef 39, Amsterdam 1105 AZ, the Netherlands

⁴ Department of Medical Microbiology and Infection Prevention, Amsterdam University Medical Centers, Meibergdreef 9, Amsterdam 1105 AZ, the Netherlands



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

incidence of the disease is likely to increase due to climate change and its consequences, such as regular floodings [8]. Pathogenic leptospires are often maintained in rodents, from where they spill over into humans and animals [7]. However, animals such as cattle, dogs or pigs may themselves act as maintenance hosts for some serovars [9]. While infections may often be asymptomatic, humans may suffer from an acute febrile illness that can potentially result in multi-organ failure [10]. Leptospirosis in swine is typically associated with reproductive disorders, such as abortion at every stage of pregnancy, decreased number of piglets per litter, the birth of runt piglets, increased weaning-to-oestrus interval, agalactia and the so-called SMEDI syndrome (stillbirth; mummification/maceration; embryonic death; infertility) [7, 9, 11, 12]. According to serological data, leptospirosis is common in swine worldwide, with seropositivity ranging from 13 to 36%, depending on the region [9, 13]. Molecular detection by PCR or even isolation of viable leptospires from clinical cases in swine is rarely successful, at least in Central Europe, a fact that might also be influenced by the submission of inappropriate sample material or quality. Thus, in clinical veterinary practice, leptospirosis is mainly diagnosed based on serological evidence, either by a fourfold rise in MAT-titer between paired sera, or – more often – by a single MAT-titer exceeding a certain threshold (usually 1:100) [9]. Since MAT-reactivity is strongly dependent on the relatedness of the infecting strain against which antibodies are developed and the test strain, poor knowledge of locally prevailing *Leptospira* species strongly hampers diagnostic conclusiveness of MAT, with a high probability of the latter to provide false negative test results, due to an incomplete panel of leptospiral serovars employed for the test [14]. Consequently, it is possible that infection with exotic species or serogroups will be overlooked, posing a potential threat to people exposed to livestock.

Leptospira (*L.*) *licerasiae* is a species of intermediate pathogenicity belonging to subclade P2 that was first detected in symptomatic human patients and peridomestic rats in urban, peri-urban, and rural areas of the Iquitos region of the Peruvian Amazon [15]. Serological evidence of infection was found in collared peccaries from the same region [16]. Later, it was also isolated from a Japanese traveler returning from Brazil [17]. Isolation of *L. licerasiae* from environmental samples collected in the Philippines and Malaysia demonstrated that the species is not restricted to South America [18, 19]. Interestingly, *L. licerasiae* was also identified as biopharmaceutical cell culture production contaminant [20]. So far, no reports have demonstrated the identification or isolation of *L. licerasiae* from Europe or animals other than those listed above.

Here, we report the first detection and genetic characterization of *L. licerasiae* isolated from a clinically healthy pig from Austria. Since leptospirosis resulting from *L. licerasiae* infection is easily overlooked by standard molecular and serological diagnostic tools, humans and animals might be exposed by a greater extent to potentially harmful leptospires than previously expected.

Case presentation

In the course of an experimental infection to characterize the pathogenesis of *L. interrogans* serovar Icterohaemorrhagiae that is described in detail elsewhere [21], pig #15 served as one of three contact animals (six months old). Contact animals were co-housed with experimentally infected pigs ($n=3$) after challenge. Control animals ($n=2$) were housed separately throughout the study. All pigs were purchased from the same source (private breeder). All pigs tested negative (MAT titers $\leq 1:50$) for the *Leptospira* serovars Icterohaemorrhagiae, Bratislava, Canicola, Grippotyphosa, Pomona, Wolffi, Tarassovi, and Hardjo before enrollment into the experiment and contact pigs were co-housed with the experimentally infected animals starting from three hours after challenge until termination of the study. Animals in the challenge group were infected with *L. interrogans* serovar Icterohaemorrhagiae. Blood, urine as well as vaginal swab samples were collected on ten time-points during the study, which was terminated 28 days post challenge by euthanasia of all experimental animals. During necropsy, liver, kidneys, urinary bladder, ovaries, oviducts, and both uterine horns were sampled as described [21]. Real-time PCR specifically targeting the leptospiral *lipL32* gene, cultivation, and MAT against the serovars Icterohaemorrhagiae, Bratislava, Canicola, Grippotyphosa, Pomona, Wolffi, Tarassovi, and Hardjo were performed as described elsewhere [21–23].

A single liquid EMJH-culture from a urine sample collected from pig #15 on day 10 post infection of the challenge group showed spirochetal growth after eight days of incubation. The ample presence of roughly 9–12 μm long, thin, helically coiled, spirochetes with hooked ends was demonstrated in this culture (termed Ages40_isolate) by electron microscopy. The bacteria consisted of a homogeneous protoplasm and two periplasmic flagella (Fig. 1). By PCR, both the urine sample and the culture were negative for the *lipL32* gene, that is associated with pathogenic *Leptospira* [23], but positive by 16S real-time PCR [24] with both probes. No sample from any of the other contact pigs or experimentally infected animals tested positive by either PCR or cultivation. To further identify the spirochetal species, present in, and isolated from the urine sample of pig #15, 16S rRNA PCR and Sanger sequencing was performed (see Supplementary




Fig. 1 Negative staining electron microscopy from the AGES40_isolate culture medium showing several tightly coiled leptospires, with two periplasmic flagella (F, insert)


Material). This revealed 620 nucleotide-long 16S rRNA sequences from both the urine sample and the Ages40_isolate that were 100% identical to several *L. licherasiae* sequences deposited in GenBank. By that it became evident that the Ages40_isolate was not the result of transmission of the challenge strain *L. interrogans* serovar Icterohaemorrhagiae, used to infect pigs in the animal experiment. Upon serotyping, the Ages40_isolate was unreactive in MAT against a panel of 43 anti-*Leptospira* rabbit reference sera and six additional rabbit antisera, including serogroup Iquitos serovar Varillal strain VAR 010 (Supplementary Table 1). However, when serum of pig#15 was tested in MAT against the AGES40_isolate and strain VAR_010, serovar Patoc strain Patoc I and serovar Andamana strain CH11, it reacted with final titres of 1:1280, 1:2560, 1:20 and 1:80, respectively.

To further elucidate the identity of the Ages40_isolate, whole genome sequencing of the isolate was performed, which revealed the presence of a second bacterium that was identified as *Brevundimonas vesicularis*, by bioinformatic analysis (Supplementary Fig. 1) and MALDI-TOF analysis. After the assembly had been cleaned from *Brevundimonas vesicularis* contigs, the final assembly resulted in a 4.28Mbp genome (N50: 138kbp, 82 contigs). We compared the newly sequenced genome to 33 genomes covering different species within the genus *Leptospira* and built a phylogenomic tree based on the

core genome of these genomes. The newly sequenced genome clearly clustered with other genomes of the *L. licherasiae* species (Fig. 2). To verify the organisms assignment to *L. licherasiae*, we constructed a pangenome and calculated the average nucleotide identity to the same 33 genomes, which confirmed our species assignment (Supplementary Figs. 2 and 3).

Discussion and conclusions

Despite frequent serological evidence, leptospiral infections in livestock are still poorly understood. Awareness about locally prevalent *Leptospira* species and serotypes, their zoonotic potential, the significance of laboratory tests and clinical presentation as well as the prevalence of subclinical infection is presently missing. Here, we report the isolation and characterization of a leptospiral species, *Leptospira licherasiae*, hitherto considered exotic to Europe, from an Austrian pig. This detection happened as a by-product of an experimental trial [21]. No influence of this result on the final conclusions of the previously published animal experiment was conceived, as the finding was limited to an unchallenged contact pig without evidence of clinical or pathological alterations and because any transmission of the challenge strain from the infected pigs was clearly excluded. A shortcoming of the work presented here is the fact that – despite 0.8/0.2 µM filtration followed by



plating on solid EMJH-agar – we were not yet able to provide a pure, uncontaminated culture of *L. licerasiae*, that would be necessary for further serological studies with MAT, i.e. to screen both animals such as kept pigs and humans with close contact to them for prior exposure. Thus, so far, infection with *L. licerasiae* has been confirmed in a single pig only. Infection of pig #15 is further supported by the MAT results, where serum from this animal strongly reacted with both the homologous isolate as well as with the related *L. licerasiae* VAR_010 strain. These titers represent a pattern of heterogeneous reactions that is usually observed in MAT, as antibodies formed during the early infection stage are not strictly specific to the homologous serovar [25, 26].

Since its first description, *L. licerasiae* has been detected by cultural isolation and/or molecular detection in several locations around the world [18, 27, 28], mostly from tropical or subtropical zones. Furthermore, outside of Peru (South America), *L. licerasiae* has so far been described only in studies of the environmental microbiome, but was never isolated from an animal or human host, except in a traveller returning from Brazil [17]. Austria, located in the temperate zone of Europe, is thus a hitherto unknown region for the occurrence of *L. licerasiae* and the fact that it was isolated from a domestic pig makes this finding even more significant. In South America, the potential reservoir animals of this organism are peridomestic rodents and members of the family Tayassuidae, including collared peccaris (*Tayassu tajacu*) [15, 16]. It is currently unclear if European pigs might play a similar role. It is, however, conceivable that *L. licerasiae* infections in Europe have been overlooked so far, as they go undetected by standard molecular and serological diagnostic tools, as was also seen in our study.

Several reports indicate a possible involvement of *L. licerasiae* in human disease [15, 17]. Detection of other members of the “intermediate pathogenic” group of leptospires (now classified as P2) in febrile human patients was documented [29]. Genomic comparisons have also argued that *L. licerasiae* is closer related to pathogenic than to saprophytic *Leptospira*, by sharing more genes with pathogenic strains and having similar metabolic traits, such as Vitamin B12 de novo biosynthesis capabilities [30]. Nevertheless, the *L. licerasiae* lipopolysaccharide composition was found to be substantially different from the one of a pathogenic *Leptospira* species [31].

The identification of *L. licerasiae* in swine from central Europe is at first surprising; however, this might just be a reflection of our patchy knowledge of the global distribution of *Leptospira* species, especially those of intermediate pathogenicity. This picture is likely to change with the ongoing collection of data from environmental studies. It is currently unknown, if

L. licerasiae is an organism commonly present in swine or in other domestic and wild animals in Europe, let alone if it causes disease in susceptible hosts.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12917-024-04213-6>.

Supplementary Material 1.
Supplementary Material 2.
Supplementary Material 3.
Supplementary Material 4.
Supplementary Material 5.
Supplementary Material 6.
Supplementary Material 7.

Acknowledgements

The excellent technical support by the Department for Molecular Biology, Institute for Veterinary Disease Control Mödling, is acknowledged.

Authors' contributions

AS performed data analysis and wrote the manuscript. AS, CU and RS conceptualized the work. DW and MS performed MAT and bacterial isolation. SR performed electron microscopy. AAA arranged MALDI-TOF and performed PCR and Sanger sequencing. HvdL performed MAT serotyping. DRM and NP performed bioinformatic analysis of NGS data. All authors reviewed the manuscript.

Funding

Not applicable.

Availability of data and materials

Next generation sequencing data of the newly described isolate was deposited at DDBJ/ENA/GenBank under the accession JBBBVK000000000. Metadata and accession numbers of the reference sequences used for construction of the phylogenomic tree, the pangenome and the nucleotide identity comparison are listed in Supplementary Table 3.

Declarations

Ethics approval and consent to participate

Ethical approval of the animal experiment, on which the presented work is partly based upon was obtained as described in Steinparzer et al., 2023 [21].

Consent for publication

Not applicable

Competing interests

The authors declare no competing interests.

Received: 20 February 2024 Accepted: 30 July 2024

Published online: 07 August 2024

References

- Korba AA, Lounici H, Kainiu M, Vincent AT, Mariet JF, Veyrier FJ, et al. *Leptospira ainlahdjerensis* sp. nov., *Leptospira ainzelensis* sp. nov. *Leptospira abararensis* sp. nov. and *Leptospira chreensis* sp. nov. four new species isolated from water sources in Algeria. *Int J Syst Evol Microbiol*. 2021;71(12):005148.
- Vincent AT, Schiettekatte O, Goarant C, Neela VK, Bernet E, Thibeaux R, et al. Revisiting the taxonomy and evolution of pathogenicity of the

- genus *Leptospira* through the prism of genomics. *PLoS Negl Trop Dis*. 2019;13(5):e0007270.
3. Perolat P, Chappel RJ, Adler B, Baranton G, Bulach DM, Billingham ML, et al. *Leptospira fainei* sp. nov., isolated from pigs in Australia. *Int J Syst Bacteriol*. 1998;48(Pt 3):851–8.
 4. Picardeau M. Virulence of the zoonotic agent of leptospirosis: still terra incognita? *Nat Rev Microbiol*. 2017;15(5):297–307.
 5. Marquez A, Djelouadi Z, Lattard V, Kodjo A. Overview of laboratory methods to diagnose Leptospirosis and to identify and to type leptospires. *Int Microbiol*. 2017;20(4):184–93.
 6. Victoriano AF, Smythe LD, Gloriani-Barzaga N, Cavinta LL, Kasai T, Limpakarnjanarat K, et al. Leptospirosis in the Asia Pacific region. *BMC Infect Dis*. 2009;9:147.
 7. Adler B, de la Pena MA. Leptospira and leptospirosis. *Vet Microbiol*. 2010;140(3–4):287–96.
 8. Picardeau M. Leptospira and Leptospirosis. *Methods Mol Biol*. 2020;2134:271–5.
 9. Strutzberg-Minder K, Tschentscher A, Beyerbach M, Homuth M, Kreienbrock L. Passive surveillance of *Leptospira* infection in swine in Germany. *Porcine Health Manag*. 2018;4:10.
 10. India W. Leptospirosis laboratory manual: WCO India; 2007. Available from: <https://iris.who.int/handle/10665/205429>.
 11. Behera SK, Sabarinath T, Mishra PKK, Deneke Y, Kumar A, ChandraSekar S, et al. Immunoinformatic study of recombinant LigA/BCon1-5 antigen and evaluation of its diagnostic potential in primary and secondary binding tests for serodiagnosis of porcine leptospirosis. *Pathogens*. 2021;10(9):1082.
 12. Pozzi P, Alborali GL, Etinger M, Hadani Y. Epidemiological investigation of the prevalence of *Leptospira* Spp. In Pigs in Israel. *Israel J Vet Med*. 2020;75(2):8.
 13. Gomes de Araujo H, Limeira CH, Ferreira Viviane, de Aquino V, Longo Ribeiro Vilela V, Jose Alves C, Dos Santos Silvano, Higinio S, et al. Global seropositivity of swine leptospirosis: systematic review and meta-analysis. *Trop Med Infect Dis*. 2023;8(3):158.
 14. Musso D, La Scola B. Laboratory diagnosis of leptospirosis: a challenge. *J Microbiol Immunol Infect*. 2013;46(4):245–52.
 15. Matthias MA, Ricaldi JN, Cespedes M, Diaz MM, Galloway RL, Saito M, et al. Human leptospirosis caused by a new, antigenically unique *Leptospira* associated with a *Rattus* species reservoir in the Peruvian Amazon. *PLoS Negl Trop Dis*. 2008;2(4):e213.
 16. Jori F, Galvez H, Mendoza P, Cespedes M, Mayor P. Monitoring of leptospirosis seroprevalence in a colony of captive collared peccaries (*Tayassu tajacu*) from the Peruvian Amazon. *Res Vet Sci*. 2009;86(3):383–7.
 17. Tsuboi M, Koizumi N, Hayakawa K, Kanagawa S, Ohmagari N, Kato Y. Imported *Leptospira icterohaemorrhagiae* Infection in Traveler Returning to Japan from Brazil. *Emerg Infect Dis*. 2017;23(3):548–9.
 18. Azali MA, Yean Yean C, Harun A, Aminuddin Baki NN, Ismail N. Molecular characterization of *Leptospira* spp. in environmental samples from North-Eastern Malaysia revealed a pathogenic strain, *Leptospira alstonii*. *J Trop Med*. 2016;2016:2060241.
 19. Saito M, Villanueva SY, Chakraborty A, Miyahara S, Segawa T, Asoh T, et al. Comparative analysis of *Leptospira* strains isolated from environmental soil and water in the Philippines and Japan. *Appl Environ Microbiol*. 2013;79(2):601–9.
 20. Chen J, Bergevin J, Kiss R, Walker G, Battistoni T, Lufburrow P, et al. Case study: a novel bacterial contamination in cell culture production-leptospira icterohaemorrhagiae. *PDA J Pharm Sci Technol*. 2012;66(6):580–91.
 21. Steinparzer R, Duerlinger S, Schmoll F, Steinrigl A, Bago Z, Willixhofer D, et al. *Leptospira interrogans* serovar icterohaemorrhagiae failed to establish distinct infection in naive gilts: lessons learned from a preliminary experimental challenge. *Pathogens*. 2023;12(1):135.
 22. Goris MG, Hartskeerl RA. Leptospirosis serodiagnosis by the microscopic agglutination test. *Curr Protoc Microbiol*. 2014;32:Unit 12E 5.
 23. Ahmed AA, Goris MGA, Meijer MC. Development of lipL32 real-time PCR combined with an internal and extraction control for pathogenic *Leptospira* detection. *PLoS ONE*. 2020;15(11):e0241584.
 24. Waggoner JJ, Balassiano I, Abeynayake J, Sahoo MK, Mohamed-Hadley A, Liu Y, et al. Sensitive real-time PCR detection of pathogenic *Leptospira* spp. and a comparison of nucleic acid amplification methods for the diagnosis of leptospirosis. *PLoS One*. 2014;9(11):e112356.
 25. Borg-Petersen C, Fagraeus A. The influence of the antigen density and other factors on the serum titer in the agglutination-lysis-test for leptospirosis. *Acta Pathol Microbiol Scand*. 1949;26(4):555–67.
 26. Kmety E. Paradox reactions and their significance in serodiagnosis of some types of leptospirosis. *Zentralbl Bakteriol Orig*. 1958;170(8):597–608.
 27. Masuzawa T, Sakakibara K, Saito M, Hidaka Y, Villanueva S, Yanagihara Y, et al. Characterization of *Leptospira* species isolated from soil collected in Japan. *Microbiol Immunol*. 2018;62(1):55–9.
 28. Sato Y, Hermawan I, Kakita T, Okano S, Imai H, Nagai H, et al. Analysis of human clinical and environmental *Leptospira* to elucidate the eco-epidemiology of leptospirosis in Yaeyama, subtropical Japan. *PLoS Negl Trop Dis*. 2022;16(3):e0010234.
 29. Chiriboga J, Barragan V, Arroyo G, Sosa A, Birdsell DN, Espana K, et al. High prevalence of intermediate *Leptospira* spp. DNA in febrile humans from Urban and Rural Ecuador. *Emerg Infect Dis*. 2015;21(12):2141–7.
 30. Ricaldi JN, Fouts DE, Selengut JD, Harkins DM, Patra KP, Moreno A, et al. Whole genome analysis of *Leptospira icterohaemorrhagiae* provides insight into leptospiral evolution and pathogenicity. *PLoS Negl Trop Dis*. 2012;6(10):e1853.
 31. Patra KP, Choudhury B, Matthias MM, Baga S, Bandyopadhyay K, Vinetz JM. Comparative analysis of lipopolysaccharides of pathogenic and intermediate pathogenic *Leptospira* species. *BMC Microbiol*. 2015;15:244.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.