## **CORRECTION**



## Correction to: Cytotoxicity of *Prymnesium parvum* extracts and prymnesin analogs on epithelial fish gill cells RTgill-W1 and the human colon cell line HCEC-1CT

Elisabeth Varga<sup>1,2</sup> • Hélène-Christine Prause<sup>1,3</sup> • Matthias Riepl<sup>1</sup> • Nadine Hochmayr<sup>1</sup> • Deniz Berk<sup>1</sup> • Eva Attakpah<sup>1</sup> • Endre Kiss<sup>1,4</sup> • Nikola Medić<sup>5,6</sup> • Giorgia Del Favero<sup>1,4</sup> • Thomas Ostenfeld Larsen<sup>7</sup> • Per Juel Hansen<sup>5</sup> • Doris Marko<sup>1</sup>

Published online: 9 April 2024 © The Author(s) 2024

## **Correction to:**

Archives of Toxicology (2024) 98:999–1014 https://doi.org/10.1007/s00204-023-03663-5

In this article, the structures of yyyy two chlorine atoms were missing in Fig. 1 and should have been appeared as: Incorrect version:

The original article can be found online at https://doi.org/10.1007/s00204-023-03663-5.

- Elisabeth Varga elisabeth.varga@vetmeduni.ac.at
- Department of Food Chemistry and Toxicology, Faculty of Chemistry, University of Vienna, Währinger Str. 38-40, 1090 Vienna, Austria
- Unit Food Hygiene and Technology, Institute of Food Safety, Food Technology and Veterinary Public Health, University of Veterinary Medicine, Vienna, Veterinärplatz 1, 1210 Vienna, Austria
- Vienna Doctoral School in Chemistry, Faculty of Chemistry, University of Vienna, Währinger Str. 42, 1090 Vienna, Austria

- Core Facility Multimodal Imaging, Faculty of Chemistry, University of Vienna, Währinger Str. 38-42, 1090 Vienna, Austria
- Marine Biological Section, Department of Biology, University of Copenhagen, Strandpromenaden 5, 3000 Helsingør, Denmark
- <sup>6</sup> Center for Bioresources, Division for Food and Production, Danish Technological Institute, Gregersensvej 8, 2630 Taastrup, Denmark
- Department of Biotechnology and Biomedicine, Technical University of Denmark, Søltofts Plads 221, 2800 Kgs Lyngby, Denmark



## Corrected version:

**Fig. 1** Backbone structure of **A**- (top) and **B**-type (bottom) prymnesins according to Igarashi et al. (1996) and Rasmussen et al. (2016a), respectively. The main difference is the length of the carbon backbone, further modifications of

prymnesin analogs are attached sugar moieties, the degree of saturation and the number of incorporated chloride and oxygen atoms

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

