

ORIGINAL ARTICLE OPEN ACCESS

Asthma and Lower Airway Disease

Methodology for the Development of the Allergic Rhinitis and Its Impact on Asthma (ARIA)-EAACI 2024–2025 Guidelines: From Evidence-to-Decision Frameworks to Digitalised Shared Decision-Making Algorithms

Jean Bousquet^{1,2,3}  | Bernardo Sousa-Pinto^{4,5} | Rafael José Vieira^{4,5}  | Holger J. Schünemann^{2,6,7} | Torsten Zuberbier^{1,2,6}  | Antonio Bognanni⁶  | Alkis Togias⁸ | Boleslaw Samolinski⁹ | Arunas Valiulis^{10,11} | Sian Williams¹² | Anna Bedbrook³ | Wienczyslaw Czarlewski^{3,13} | Maria Jose Torres¹⁴  | Mohamed H. Shamji¹⁵  | Mário Morais-Almeida¹⁶  | G. Walter Canonica^{7,17}  | Leticia de las Vecillas¹⁸  | Mark S. Dykewicz¹⁹ | Cristina Jacomelli²⁰ | Ludger Klimek^{21,22}  | Lucas Leemann²³ | Olga Lourenço²⁴  | Yuliia Palamarchuk²⁵ | Nikolaos G. Papadopoulos²⁶  | Ana Margarida Pereira^{4,27,28} | Marine Savoure^{29,30,31}  | Sanna K. Toppila-Salmi^{32,33}  | Maria Teresa Ventura^{34,35} | Juan Jose Yepes-Nuñez^{36,37} | Alvaro A. Cruz³⁸  | Giorgio Ciprandi³⁹ | Bilun Gemicioglu⁴⁰ | Mattia Giovannini^{41,42}  | Brigita Gradauskiene⁴³ | Tuomas Jartti⁴⁴  | Miloš Jeseňák^{45,46,47}  | Piotr Kuna⁴⁸ | Violeta Kvedariene^{49,50} | Désirée E. Larenas-Linnemann⁵¹ | Amir H. A. Latiff⁵² | Yousser Mohammad^{53,54} | Ken Ohta⁵⁵ | Padukudru A. Mahesh⁵⁶ | Isabella Pali-Schöll⁵⁷ | Oliver Pfaar⁵⁸  | Frederico S. Regateiro^{59,60,61,62} | Nicolas Roche^{63,64,65} | Mikhail Sofiev²⁵ | Luís Taborda-Barata^{24,66,67} | Charlotte Suppli Ulrik^{68,69} | Marylin Valentin Rostan⁷⁰ | Giovanni Viegi⁷¹ | Luo Zhang⁷²  | Josep M. Antó^{30,31,73} | Tari Haahtela⁷⁴  | Ivan Cherrez-Ojeda^{1,2,75,76}  | Juan Carlos Ivancevich⁷⁷ | Nikolai Khaltayev⁷⁸ | Arzu Yorgancioglu⁷⁹  | Baharudin Abdullah⁸⁰ | Mona Al-Ahmad⁸¹ | Maryam Ali Al-Nesf⁸²  | Rita Amaral^{14,5,83,84}  | Julijana Asllani⁸⁵  | Karl-C Bergmann^{1,2} | Jonathan A. Bernstein⁸⁶  | Michael S. Blaiss⁸⁷ | Fulvio Baido^{88,89} | Paulo Camargos⁹⁰ | Pedro Carreiro-Martins^{91,92} | Thomas Casale⁹³ | Lorenzo Cecchi⁹⁴  | Alessandro G. Fiocchi⁹⁵  | Antonio F. M. Giuliano⁹⁶  | George Christoff⁹⁷ | Ieva Cirule⁹⁸ | Jaime Correia-de-Sousa⁹⁹  | Elisio M. Costa¹⁰⁰ | Stefano Del Giacco¹⁰¹  | Philippe Devillier¹⁰² | Dejan Dokic¹⁰³  | Elham Hosny¹⁰⁴ | Tomohisa Iinuma¹⁰⁵  | Carla Irani¹⁰⁶ | Zhanat Ispayeva¹⁰⁷ | Kaja Julge¹⁰⁸ | Igor Kaidashev¹⁰⁹ | Kazi S. Bennoor¹¹⁰ | Helga Kraxner¹¹¹ | Inger Kull^{112,113}  | Marek Kulus¹¹⁴ | Maciej Kupczyk⁴⁶ | Andriy Kurchenko¹¹⁵  | Stefania La Grutta¹¹⁶ | Neven Miculinic¹¹⁷ | Lan Le Thi Tuyet¹¹⁸ | Michael Makris¹¹⁹ | Branislava Milenkovic¹²⁰ | Sang Min Lee¹²¹  | Stephen Montefort¹²² | André Moreira^{123,124,125} | Joaquim Mullol¹²⁶ | Rachel Nadir^{127,128}  | Alla Nakonechna^{129,130} | Hugo E. Neffen¹³¹ | Marek Niedozytko¹³²  | Dieudonné Nyembue¹³³ | Robyn O'Hehir¹³⁴  | Ismail Ogulur¹³⁵ | Yoshitaka Okamoto^{136,137} | Heidi Olze¹³⁸ | Oscar Palomares¹³⁹  | Petr Panzner¹⁴⁰  | Vincenzo Patella^{141,142,143}  | Ruby Pawankar¹⁴⁴  | Constantinos Pitsios¹⁴⁵ | Todor A. Popov¹⁴⁶ | Francesca Puggioni¹⁴⁷ | Santiago Quirce¹⁸ | Agné Ramonaitė^{148,149} | Marysia Recto¹⁵⁰ | Maria Susana Repka-Ramirez¹⁵¹  | Graham Roberts^{152,153,154}  | Karla Robles-Velasco^{31,74,155}  | Menachem Rottem^{156,157} | Marianella Salapatras¹⁵⁸ | Joaquin Sastre¹⁵⁹ | Nicola Scichilone¹⁶⁰ | Juan Carlos Sisul¹⁶¹ | Dirceu Solé¹⁶² | Manuel E. Soto-Martinez¹⁶³ | Milan Sova¹⁶⁴ | Pongsakorn Tantilipikorn¹⁶⁵ | Ana Todo-Bom¹⁶⁶ | Vladyslav Tsaryk¹¹³  | Ioanna Tsiligianni^{167,168} | Marilyn Urrutia-Pereira¹⁶⁹ | Erkka Valovirta¹⁷⁰ | Eric Van Ganse¹⁷¹ | Tuula Vasankari^{172,173} | Dana Wallace¹⁷⁴ | De Yun Wang¹⁷⁵  | Margitta Worm¹⁷⁶  | Osman M. Yusuf¹⁷⁷  | Mihaela Zidarn^{178,179} | Sara Gil-Mata^{4,5} | Manuel Marques-Cruz^{4,5} | Bassam Mahboub^{180,181} | Ignacio J. Ansotegui¹⁸² | Antonino Rouhard^{183,184} | Maria Cristina Artesani⁹⁵  | Bruno Barreto¹⁸⁵ | Sven Becker¹⁸⁶  | Bianca Beghe¹⁸⁷ | Jacques Bouchard¹⁸⁸ | Melisande Bourgoin-Heck^{189,190,191}  | Luisa Brussino^{192,193} | Roland Buhl¹⁹⁴ | Mario Calvo-Gil¹⁹⁵ | Victoria Cardona Dahl^{196,197} | José Antonio Castillo Vizuete^{198,199} | Denis Charpin²⁰⁰ | Niels H. Chavannes²⁰¹ | Marta Chełmińska²⁰² | Lei Cheng²⁰³ | Ekaterine Chkhartishvili²⁰⁴ | Herberto Jose Chong-Neto²⁰⁵ | Deepa Choudhury²⁰⁶ | Derek Chu^{207,208} | Cemal Cingi²⁰⁹  | Enrico Compalati²¹⁰ | Biljana Cvetkovski²¹¹ | Jane Da Silva^{212,213} | Gennaro D'Amato²¹⁴  | Janet Davies^{215,216} | Danilo Di Bona²¹⁷ | Maria V. Dimou²⁶ | Maria Do Ceu Teixeira^{218,219} | Maria Doulaptsi²²⁰  | José Miguel Fuentes Pérez²²¹  | Radosław Gawlik²²² | Ozlem Goksel²²³  | Maximiliano R. Gómez²²⁴ |

Dr. Alkis Togias' co-authorship of this report does not constitute endorsement by the National Institute of Allergy and Infectious Diseases, the National Institutes of Health, or any other agency of the United States Government.

For affiliations refer to page 445.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2025 The Author(s). *Allergy* published by European Academy of Allergy and Clinical Immunology and John Wiley & Sons Ltd.

Sandra N. Gonzalez Diaz²²⁵  | Maia Gotua²²⁶ | Christos Grigoreas²²⁷ | Ineta Grisle²²⁸ | Maria Antonieta Guzman²²⁹ | Rachel House Tan^{230,231} | Michael Hyland²³² | Despo Ierodiakonou²³³ | Aspasia Karavelia²³⁴ | Marta Kisiel²³⁵ | Mitja Kosnik^{178,179}  | Vicky Kritikos^{236,237}  | Carlo Lombardi²³⁸ | Matteo Martini^{239,240}  | Cem Meço^{241,242,243} | Eris Mesonjesi²⁴⁴ | Florin Mihaltan²⁴⁵ | Marcin Moniuszko²⁴⁶ | Robert Naclerio²⁴⁷ | Sophia Neisinger^{1,2} | Michal Ordak²⁴⁸  | Giovanni Paoletti^{7,147}  | Edgar Arturo Perdomo-Flores²⁴⁹ | Nhan Pham-Thi^{250,251,252} | Emmanuel Prokopakis²²⁰ | Daniela Rivero Yeverino²⁵³ | Giovanni Rolla¹⁹² | Jan Romantowski¹³² | Philip W. Rouadi^{254,255}  | Maia Rukhadze²⁵⁶ | Daiju Sakurai²⁵⁷ | Laila Salameh²⁵⁸ | Faradiba Serpa Sarquis²⁵⁹ | Tanja Soklic Kosak²⁶⁰  | Michael Soyka²⁶¹  | Olga Sozinova²⁶²  | Krzysztof Specjalski¹³² | Vesna Tomic-Spiric^{263,264} | Martina Vachova^{265,266} | Marianne van Hage^{267,268,269}  | Ilgim Vardaloglu Koyuncu²⁷⁰ | Pakit Vichyanond^{271,272} | Martin Wagenmann²⁷³  | Fanny Wai San Ko²⁷⁴ | Pascal Werminghaus²⁷⁵ | Vicky Paraskevi Xepapadaki²⁶ | Yi-Kui Xiang^{1,2,276}  | João A. Fonseca^{4,5}

Correspondence: Jean Bousquet (jean.bousquet@orange.fr)

Received: 17 July 2025 | **Accepted:** 3 August 2025

Funding: FRAUNHOFER Institute for Translational Medicine and Pharmacology ITMP, Immunology and Allergology, Berlin, Germany; ARIA (Allergic Rhinitis and its Impact on Asthma), Montpellier, France; CINTESIS@RISE, Porto, Portugal.

Keywords: allergic rhinitis | asthma | ENT (rhinitis, sinusitis, nasal polyps...) | guidelines

ABSTRACT

The Allergic Rhinitis and its Impact on Asthma (ARIA) guidelines produced their first edition in 1999, with subsequent revisions in 2008, 2010, 2016 and 2019. A new iteration of ARIA—ARIA 2024–2025—in collaboration with EAACI is currently being developed, focusing on the management of allergic rhinitis. ARIA 2024–2025 follows the GRADE framework and is endorsed by the European Academy of Allergy and Clinical Immunology (EAACI). A set of approaches has been used to develop guideline questions, including surveying key opinion leaders and using artificial intelligence (AI)-based tools to analyse web searches on allergic rhinitis and to generate questions. Each prioritised guideline question is assessed through an Evidence-to-Decision (EtD) framework. EtDs support the systematic and transparent formulation of recommendations, comprising 12 criteria for which the best available evidence should be sought. In the context of ARIA-EAACI 2024–2025, such evidence is derived not only from randomised controlled trials but also—among others—from patient-generated data sources that better reflect the affected individuals' perspectives. Moreover, ARIA-EAACI 2024–2025 incorporates evidence on planetary health. Developed guideline recommendations will support the creation of digitalised decision algorithms and care pathways. This paper describes the methodology used to develop the person-centred, digitally enabled and AI-assisted ARIA-EAACI 2024–2025. Among others, it describes (i) the development and prioritisation of guideline questions, (ii) sources of evidence for EtDs and (iii) the development of digitalised decision algorithms and care pathways.

1 | Introduction

Allergic rhinitis (AR) is one of the most common chronic conditions globally, often co-occurring with asthma and conjunctivitis [1–3]. It impairs quality of life and affects social life, school and work productivity, and is associated with substantial economic costs [1, 4]. AR is an excellent candidate for testing innovation in guidelines: the vast inter-individual differences in exposures and in personal beliefs, preferences and values require a person-centred approach. In addition, AR is often self-managed, with limited contact with health services.

The Allergic Rhinitis and its Impact on Asthma (ARIA) initiative first proposed guidelines for AR and asthma multimorbidity in 1999 [5]. ARIA has evolved from the first multimorbidity guideline in respiratory diseases using the Shekelle evidence-based model [5] to the GRADE (Grading of Recommendations, Assessment, Development and Evaluation) framework for evidence-based guidelines [6, 7]. In fact, ARIA 2016 was used as the clinical scenario on how to interpret and use a guideline in a review published in the *Journal of the American Medical Association* [8]. ARIA 2010 and 2016 devised a medication algorithm [6, 7]. However, digitalisation was not implemented.

In 2019, next-generation guidelines using real-life MASK-air data and allergen exposure chamber studies proposed digitally enabled person-centred care for the first time [9]. The next iteration, ARIA-EAACI 2024–2025, is endorsed by the European Academy of Allergy and Clinical Immunology and will result in a person-centred, digitally enabled, artificial intelligence (AI)-assisted guideline [10] (Figure 1).

ARIA-EAACI 2024–2025 is formulating recommendations based on the use of the GRADE Evidence-to-Decision (EtD) framework. The EtD supports the systematic and transparent formulation of recommendations for each prioritised guideline question [11, 12]. It comprises 12 criteria for which the best available evidence should be sought. In ARIA-EAACI 2024–2025, such evidence is not only based on data from randomised controlled trials (RCTs) but also from other sources, including studies using data directly provided by patients. In fact, although RCTs are the gold standard for assessing the efficacy of medical interventions, they present several limitations. In this context, in AR, well-designed pragmatic trials and observational studies using direct patient data can be a source of complementary evidence (Bousquet, submitted), providing insights into the reality of day-to-day clinical practice.




FIGURE 1 | Graphical representation of the tasks of the Allergic Rhinitis and its Impact on Asthma (ARIA-EAACI) 2024–2025 guidelines and their classification as patient-centred, digitally enabled and artificial intelligence (AI)-assisted.

In addition to being informed by multiple sources of evidence, ARIA-EAACI 2024–2025 will be innovative in its final result (Table 1). In addition to recommendations, ARIA-EAACI 2024–2025 will result in a set of personalised and digitalised algorithms that can be embedded in an app. For example, the MASK-air mHealth app is a Class IIa medical device, in which algorithms can be tested and implemented into the clinical workflow of allergists [13, 15]. Henceforth, recommendations will be included in digitalised management algorithms considering multiple clinical scenarios. Algorithm digitalisation has several advantages, including patient empowerment and flexibility for the implementation of updates in the perspective of a living ARIA guideline. Digital divide (e.g., age-related) aspects do however need to be considered.

In this paper, we aim to present the methodology for the development of ARIA 2024–2025 and to provide a rationale for the creation of digitalised guideline algorithms.

2 | ARIA-EAACI 2024–2025: Patient-Centred, Digitally Enabled, AI-Assisted Guidelines

2.1 | Patient Empowerment

ARIA-EAACI 2024–2025 is being devised throughout the different stages on a person-centred basis (Figure 1). First, AR patients have been included in the guideline panel. Second, to ensure that guideline questions are person-centred, questions were formulated pertaining to (i) findings and hypotheses from studies based on data directly provided by patients

in MASK-air and (ii) AI-supported approaches that analysed popular online search queries on AR posed by internet users [16].

Third, we are using the EtD framework to develop our recommendations—the EtDs imply that the formulation of recommendations is not solely based on the desirable and undesirable effects of the interventions, but also on aspects such as patients' values and the acceptability of interventions by the different interest holders (considering aspects such as patients' satisfaction with medications or the speed of onset of action of treatments) [11, 12].

Fourth, patient behaviours are being considered to support the formulation of recommendations. As an example, we are considering steroid phobia for recommendations related to intranasal corticosteroids (INCS) [17]. In addition, we are considering that most patients do not use medication regularly, but rather on an as-needed basis. This has been observed in studies using MASK-air direct patient data [18], and the few RCTs comparing as-needed versus regular treatment for AR did not find any major differences [19–26].

Finally, the ARIA-EAACI 2024–2025 guidelines will result in the development of management algorithms that will be implemented in an mHealth app that can be accessed by patients.

2.2 | Digital Enablement

Digital enablement is a cornerstone of ARIA-EAACI 2024–2025: (i) findings and hypotheses from studies using mHealth data

TABLE 1 | Summary of innovative approaches followed in the development of the ARIA-EAACI 2024–2025 guidelines.

Task	“Classical” approaches followed in the ARIA-EAACI 2024–2025 guidelines	Innovative approaches followed in the ARIA-EAACI 2024–2025 guidelines
Formulation of guideline questions	– Identification of previous guideline questions; – Formulation of guideline questions by experts.	– Identification of questions based on direct patient-reported data (MASK-air app); – AI-assisted identification of questions posed by Internet users in online searches; – Direct generation of questions by AI tools.
Identification of outcomes	– Identification of outcomes by experts.	– Identification of outcomes by AI tools.
Completing the evidence-to-decision frameworks		
Desirable and undesirable effects	– Use of evidence from systematic reviews (of RCTs).	– Use of pharmacovigilance data.
Values	– Use of evidence from systematic reviews of values and preferences.	– Assessment of utilities based on direct patient data (MASK-air app) [study included in the systematic review of values and preferences].
Resources used and cost-effectiveness	– Use of data from the scientific literature and health technology assessment reports.	– Assessment of indirect costs based on direct patient data (MASK-air app) – Survey of ARIA experts on the costs of medications.
Equity		– Survey of ARIA experts on the availability of medication; – Systematic consideration of medications in the World Health Organisation List of Essential Medicines.
Acceptability	– Use of data from the scientific literature (e.g., for onset of action).	– Use of direct patient data (MASK-air app) to assess adherence, frequency of co-medication and treatment satisfaction with different medications.
Planetary Health	<i>(planetary health not usually considered in guidelines)</i>	– Consideration of planetary health.
Considering multimorbidity		– Explicit consideration of multimorbidity in subgroup considerations.
Building treatment algorithms	– Already proposed in the former ARIA iterations [13, 14]	– Building and implementation of digitalised algorithms in an AI-assisted process involving a multidisciplinary team.

Abbreviations: AI, artificial intelligence; RCT, randomised controlled trial.

enabled the formulation of guideline questions, (ii) mHealth direct patient data are used as a source of evidence and (iii) management algorithms will be digitalised into an mHealth app (Figure 1). A market research study has analysed available mHealth apps for AR and has concluded that MASK-air is available in the largest number of countries and has the highest number of scientific publications [27]. Moreover, MASK-air is one of the few apps reporting asthma multimorbidity. Therefore, MASK-air was selected as the app from which data are retrieved to provide evidence for guideline development and to implement digitalised algorithms.

MASK-air is available in 30 countries. It has been used by more than 40,000 patients, who have reported over 700,000 days of use [15]. MASK-air is a patient-centred Medical Device regulation Class IIa. It is a Good Practice of the Directorate General Health and Food Safety of the European Commission [28] as well as an OECD (Organisation of Economic Cooperation and

Development) Best Practice for Integrating Care to Prevent and Manage Chronic Diseases [13].

MASK-air comprises a daily monitoring questionnaire assessing the impact of rhinitis and asthma symptoms by means of validated visual analogue scales (VASs) [29, 30]. In addition, users are asked (i) to enter their daily medications via a regularly updated scroll list that contains country-specific prescribed and over-the-counter medications and (ii) to provide feedback on their satisfaction with their medication. MASK-air also includes additional questionnaires, namely the Control of Allergic Rhinitis and Asthma Test (CARAT) [31], the Work Productivity and Activity Impairment Questionnaire: Allergy Specific (WPAI-AS) [32, 33], and EQ-5D [34, 35]. Model predictions of pollen concentrations and air pollution obtained from the Finnish Meteorological Institute (<https://silam.fmi.fi>) [36, 37] and, for some species, from Copernicus Atmosphere Monitoring Service

(<http://atmosphere.copernicus.eu>) are available daily within a 10 km radius in Europe in geolocalised patients. Table S1 presents some of the relevant findings of MASK-air studies for the development of ARIA-EAACI 2024–2025.

2.3 | Use of AI

In the context of ARIA-EAACI 2024–2025, AI helped in the development of guideline questions and the identification of outcomes (Figure 1). In particular, a large language model-based chatbot was prompted to (i) classify online search queries into those conveying questions [16], (ii) suggest guideline questions [16] and (iii) suggest potentially relevant outcomes. In addition, AI was used to support the systematic review on patients' values and preferences in AR [38]. In ARIA-EAACI 2024–2025, we plan to use AI to support (i) the writing of plain language summaries and (ii) the development of the digitalised algorithms.

3 | Panel Selection

ARIA group members involved in ARIA-EAACI 2024–2025 include (i) the steering committee, (ii) the guideline panel, (iii) the ARIA review group and (iv) the ARIA Junior Members (Table S2).

The ARIA guideline panel includes 30 experts and is responsible for (i) suggesting and prioritising guideline questions and outcomes, (ii) participating in group meetings, (iii) providing input when discussing the available evidence for each guideline question, (iv) reviewing evidence summaries, (v) making judgements for each EtD criterion, (vi) drafting guideline recommendations, (vii) reviewing and writing the final report and (viii) supporting guideline dissemination.

The guideline panel was set in such a way as to ensure (i) the inclusion of experts with different profiles (physicians specialised in allergy, otorhinolaryngology, paediatrics and primary care, pharmacists, patients, methodologists and AI experts), (ii) representativeness in terms of gender, age and country and (iii) that at least 50% of the members did not have a conflict of interest (CoI). All members were required to complete the "Guideline Group or Panel Member Certification Course" (now termed "Certified Guideline Panel Member" course) of the International Guideline Training and Certification Programme INGUIDE.

To increase the diversity of the received inputs, we invited an ARIA review group (including members from low- and middle-income countries) and ARIA Junior Members to participate in certain guideline tasks. Tasks for these groups include (i) providing input on non-prioritised questions, (ii) reviewing formulated recommendations and (iii) participating in the development of the management algorithms.

4 | Development and Prioritisation of the Guideline Questions

In ARIA-EAACI 2024–2025, several different approaches were used for the development of guideline questions (Figure 2;

Table 2), including person-centred approaches. The developed questions were then prioritised to select those for which recommendations would be drafted.

4.1 | Development of Targeted Guideline Questions

4.1.1 | Building on Established Knowledge: Questions From Previous ARIA Guidelines (Physician-Centred Approach)

We retrieved all questions regarding the pharmacological and non-pharmacological treatment of AR that had been answered in ARIA 2010 and 2016 and the US Practice Parameters [39].

4.1.2 | Insights From Experts: Questions From Panel Members (Key Opinion Leader-Centred Approach)

We surveyed guideline panel members on relevant questions regarding the pharmacological or non-pharmacological treatment of AR. To avoid duplicate questions, we provided panel members with the list of guideline questions that had been answered in the previous ARIA guidelines or identified through analysis of MASK-air studies.

4.1.3 | Leveraging Direct Patient-Reported Data: Questions From MASK-air Studies (Person-Centred Approach)

To formulate questions based on observations of day-to-day practice and patients' experience, two guideline methodologists (BSP and RJV) (i) read all the original studies that used MASK-air data and (ii) formulated guideline questions based on the main messages and hypotheses raised by these studies.

4.1.4 | AI-Supported Development of Guideline Questions (Person-Centred Approach)

To better reflect patients' beliefs and needs, AI was used in the development of guideline questions [16]. We retrieved popular queries on AR (identified using Google Trends) and used ChatGPT 4.0 to classify them into those conveying potentially relevant questions or not [16]. Queries identified as potentially conveying relevant questions were then manually transformed into guideline questions.

We also prompted ChatGPT 4.0 to assume the role of a patient or of a healthcare provider and to generate potentially relevant guideline questions [16].

4.2 | Prioritisation of Guideline Questions

Developed guideline questions were voted for prioritisation by the guideline panel members who rated the priority of each question on a scale of 1–9 (9 indicating highest priority) [40] according to the criteria and signalling questions displayed in




FIGURE 2 | Flow diagram for the formulation of guideline questions. GT, Google Trends; KOL, key opinion leaders.

Box S1. For a question to be prioritised, a mean priority rate of at least 6.3 or a median rate of at least 7 was required.

Among the 89 unique developed questions, 39 were voted by the guideline panel members as “prioritised questions”. Non-prioritised questions were sent to the ARIA review group. The review group could recommend prioritising some of those questions or even propose new questions of interest. Overall, three additional questions were added further to feedback from the review group.

5 | Identification and Prioritisation of the Outcomes

For the formulation of guideline recommendations, interventions need to be judged in terms of their desirable and undesirable effects. This implies assessing the effect of interventions on a set of pre-selected outcomes (ideally not more than seven).

In ARIA-EAACI 2024–2025, a set of potentially relevant outcomes was identified by considering (i) an initial proposal by the guideline co-chairs, (ii) outputs from a large language model, (iii) subsequent suggestions by guideline panel members and (iv) a systematic review of patients’ values. For each identified outcome, we developed health outcome descriptors to create common definitions that describe the outcomes with respect to symptoms, time horizon, testing and treatment and consequences [41] (example in Box S2). This ensured that panel members displayed a common understanding when discussing each of the outcomes.

Guideline panel members were then asked to rate the priority of each outcome on a scale of 1–9 (9 indicating the highest priority). Of the 28 outcomes voted for prioritisation, the five rated as being of highest priority were nasal symptoms, quality-of-life impairment, eye symptoms, total symptoms and serious adverse events (AEs). Since these top outcomes only included one undesirable effect (serious AEs), we considered a sixth outcome

TABLE 2 | Full list of guideline questions voted for prioritisation and methods used for their generation.

Guideline question	Method for question generation				Prioritised
	ARIA 2010–2016	Guideline panel/KOL	MASK-air data	Generation based on web search queries	Direct generation by LLM
Should continuous long-term treatment vs. as-needed treatment be used in patients with allergic rhinitis?			×	×	×
Should co-medication vs. medication up dosing be used in patients with allergic rhinitis that is poorly controlled despite pharmacologic treatment?			×		×
Should pharmacologic treatment adjustment according to seasonality in pollen or other allergen exposure vs. no adjustment be used in patients with allergic rhinitis?			×		×
Should pharmacologic treatment adjustment according to air pollution levels vs. no adjustment be used in patients with allergic rhinitis?			×		×
Should intranasal H1 antihistamines vs. no treatment be used for the treatment of allergic rhinitis?	×			×	×
Should oral H1 antihistamines vs. no treatment be used for the treatment of allergic rhinitis?	×			×	×
Should intraocular H1 antihistamines vs. no treatment be used for the treatment of ocular symptoms in patients with allergic rhinitis?	×			×	×
Should intranasal glucocorticosteroids vs. no treatment be used for the treatment of allergic rhinitis?	×			×	×
Should oral leukotriene-receptor antagonists vs. no treatment be used for the treatment of allergic rhinitis?	×			×	×
Should intranasal decongestants vs. no treatment be used for the treatment of allergic rhinitis?	×			×	×
Should new-generation oral H1 antihistamines vs. old-generation oral H1 antihistamines be used for the treatment of allergic rhinitis?	×				×
Should intranasal H1 antihistamines vs. intranasal glucocorticosteroids be used for the treatment of allergic rhinitis?	×			×	×

(Continues)

TABLE 2 | (Continued)

Guideline question	Method for question generation					Prioritised
	ARIA 2010–2016	Guideline panel/KOL	MASK-air data	Generation based on web search queries	Direct generation by LLM	
Should intranasal H1 antihistamines vs. oral H1 antihistamines be used for the treatment of allergic rhinitis?	×					×
Should intranasal H1 antihistamines vs. intranasal cromones be used for the treatment of allergic rhinitis?	×					×
Should intranasal glucocorticosteroids vs. oral H1 antihistamines be used for the treatment of allergic rhinitis?	×				×	×
Should a combination of an intranasal H1 antihistamine and an intranasal glucocorticosteroid vs. an intranasal glucocorticosteroid alone be used for the treatment of allergic rhinitis?	×					×
Should a combination of an intranasal H1 antihistamine and an intranasal glucocorticosteroid vs. an intranasal H1 antihistamine alone be used for the treatment of allergic rhinitis?	×					×
Should a combination of an oral H1 antihistamine and an intranasal glucocorticosteroid vs. an intranasal glucocorticosteroid alone be used for the treatment of allergic rhinitis?	×				×	×
Should a combination of an oral H1 antihistamine and an intranasal glucocorticosteroid vs. an oral H1 antihistamine alone be used for the treatment of allergic rhinitis?		×				×
Should a combination of an oral leukotriene-receptor antagonist and an oral H1 antihistamine vs. an oral H1 antihistamine alone be used for the treatment of allergic rhinitis?		×				×
Should any specific individual intranasal glucocorticosteroid vs. other individual intranasal glucocorticosteroids be used for the treatment of allergic rhinitis?		×				×
Should a monoclonal antibody vs. no treatment be used for the treatment of patients with moderate to severe allergic rhinitis?		×				×
Should masks vs. no masks be used for preventing exposure to air pollution and other triggers in patients with allergic rhinitis?		×				×
Should air purifiers vs. no air purifiers be used for the management of patients with allergic rhinitis?		×			×	×

(Continues)

TABLE 2 | (Continued)

Guideline question	Method for question generation					Prioritised
	ARIA 2016–2016	Guideline panel/KOL	MASK-air data	Generation based on web search queries	Direct generation by LLM	
Should patient diaries vs. no patient diaries be used for monitoring in patients with allergic rhinitis?		×				×
Should digital patient diaries vs. nondigital patient diaries be used for monitoring elderly patients with allergic rhinitis?			×			×
Should a dual approach combining both long-term and short-term monitoring vs. long-term monitoring alone be used in patients with allergic rhinitis?			×			×
Should a systematic diagnostic screening for asthma vs. no systematic screening be used in patients with allergic rhinitis?			×		×	×
Should a systematic diagnostic screening for allergic rhinitis vs. no systematic screening be used in patients with asthma?			×			×
Should a systematic assessment of the effect of allergic rhinitis on quality of life, work productivity, and/or school performance vs. no systematic assessment be used in patients with allergic rhinitis?			×			×
Should therapy adherence monitoring vs. no therapy adherence monitoring be used in treated patients with allergic rhinitis?		×				×
Should modern methods to increase adherence (e.g., shared decision-making, guided self-management) vs. traditional methods to increase adherence be used for the management of patients with allergic rhinitis?		×				×
Should patient education and self-management strategies vs. no patient education and self-management strategies be used in patients with allergic rhinitis?		×			×	×
Should any specific individual oral H1 antihistamine vs. other individual oral H1 antihistamines be used for the treatment of allergic rhinitis?				×		×
Should a combination of an intranasal H1 antihistamine and an intranasal glucocorticosteroid vs. no treatment be used for the treatment of allergic rhinitis?				×		×

(Continues)

TABLE 2 | (Continued)

Guideline question	Method for question generation					Prioritised
	ARIA 2010–2016	Guideline panel/KOL	MASK-air data	Generation based on web search queries	Direct generation by LLM	
Should intraocular H1 antihistamines be used for the treatment of ocular symptoms in patients with allergic rhinitis?					×	×
Should a systematic diagnostic screening for nonallergic triggers (such as irritants or pollutants) vs. no systematic screening be used in patients with allergic rhinitis?					×	×
Should a systematic diagnostic screening for sinusitis vs. no systematic screening be used in patients with allergic rhinitis?					×	×
Should digital platforms (like smart watches, fitness trackers and mobile apps) vs. no digital platforms be used in patients with allergic rhinitis?					×	×
Should oral glucocorticosteroids vs. no treatment be used for the treatment of allergic rhinitis?	×			×		
Should intramuscular glucocorticosteroids vs. no treatment be used for the treatment of allergic rhinitis?	×					
Should intranasal chromones vs. no treatment be used for the treatment of allergic rhinitis?	×					
Should intraocular chromones vs. no treatment be used for the treatment of ocular symptoms in patients with allergic rhinitis?	×					
Should intranasal ipratropium bromide vs. no treatment be used for the treatment of allergic rhinitis?	×			×	×	
Should oral decongestants vs. no treatment be used for the treatment of allergic rhinitis?	×			×		
Should intranasal glucocorticosteroids vs. oral leukotriene-receptor antagonists be used for the treatment of allergic rhinitis?	×				×	
Should oral leukotriene-receptor antagonists vs. oral H1 antihistamines be used for the treatment of allergic rhinitis?	×				×	
Should oral leukotriene-receptor antagonists vs. intranasal H1 antihistamines be used in patients with allergic rhinitis?		×			×	

(Continues)

TABLE 2 | (Continued)

Guideline question	Method for question generation					Prioritised
	ARIA 2016–2019	Guideline panel/KOL	MASK-air data	Generation based on web search queries	Direct generation by LLM	
Should intraocular H1 antihistamines vs. nasal H1 antihistamines be used for the treatment of ocular symptoms in patients with allergic rhinitis?		×				
Should a combination of an intranasal decongestant and an oral H1 antihistamine vs. an oral H1 antihistamine alone be used for the treatment of allergic rhinitis?		×				
Should a combination of oral decongestants and H1 antihistamines vs. oral H1 antihistamines alone be used for the treatment of allergic rhinitis?	×			×		
Should nasal irrigation alone vs. no treatment be used for the treatment of allergic rhinitis?			×	×	×	
Should nasal irrigation as an add-on to pharmacologic treatment vs. pharmacologic treatment alone be used for the treatment of allergic rhinitis?			×			
Should nasal barriers (e.g., petroleum jelly) vs. no nasal barriers be used for preventing exposure to air pollution in patients with allergic rhinitis?		×				
Should sunglasses vs. no sunglasses be used for preventing exposure to air pollution and other triggers in patients with allergic rhinitis?		×				
Should temperature-controlled laminar airflow devices vs. no temperature-controlled laminar airflow devices be used for the management of patients with allergic rhinitis?		×				
Should drying bed linen and clothes indoors vs. drying bed linen and clothes outdoors be used for preventing exposure to pollen in patients with allergic rhinitis?		×			×	
Should anti-Fel d1 cat food vs. regular cat food be used in cats of patients with allergic rhinitis with allergies to cats?			×			
Should early warning systems vs. no early warning systems be used in patients with allergic rhinitis?		×				

(Continues)

TABLE 2 | (Continued)

Guideline question	Method for question generation				Prioritised
	ARIA 2010–2016	Guideline panel/KOL	MASK-air data	Generation based on web search queries	Direct generation by LLM
Should a routine olfactory test vs. no routine olfactory test be used for the monitoring of patients with allergic rhinitis?		×			
Should a systematic assessment of disease control vs. a systematic assessment of disease severity be used in patients with allergic rhinitis?		×			
Should a systematic diagnostic screening for conjunctivitis vs. no systematic screening be used in patients with allergic rhinitis?		×			
Should a systematic screening for local allergic rhinitis vs. no systematic screening for local allergic rhinitis be used in patients with rhinitis clinically triggered by airborne allergens but with negative results on skin prick tests and specific IgE to environmental allergens?		×			
Should pharmacologic treatment in the morning vs. pharmacologic treatment in the evening be used for the treatment of allergic rhinitis?				×	
Should intraocular decongestants vs. no treatment be used for the treatment of ocular symptoms in patients with allergic rhinitis?				×	
Should more than 1 daily oral H1 antihistamine vs. 1 single daily oral H1 antihistamine be used for the treatment of allergic rhinitis?				×	
Should H1 antihistamines vs. decongestants be used for the treatment of allergic rhinitis?				×	
Should any specific individual intranasal decongestant vs. other individual intranasal decongestants be used for the treatment of allergic rhinitis?				×	×
Should any specific individual oral decongestant vs. other individual oral decongestants be used for the treatment of allergic rhinitis?				×	
Should any specific individual intranasal H1 antihistamine vs. other individual intranasal H1 antihistamines be used for the treatment of allergic rhinitis?				×	

(Continues)

TABLE 2 | (Continued)

Guideline question	Method for question generation				Prioritised
	ARIA 2016–2016	Guideline panel/KOL	MASK-air data	Generation based on web search queries	Direct generation by LLM
Should any specific individual intraocular H1 antihistamine vs. other individual intraocular H1 antihistamines be used for the treatment of ocular symptoms in patients with allergic rhinitis?				×	
Should a combination of oral decongestants and acetaminophen vs. oral decongestants alone be used for the treatment of allergic rhinitis?				×	
Should a combination of oral H1 antihistamines and acetaminophen vs. oral H1 antihistamines alone be used for the treatment of allergic rhinitis?				×	
Should a combination of oral H1 antihistamines and expectorant medications (mucoactive agents) vs. oral H1 antihistamines alone be used for the treatment of allergic rhinitis?				×	
Should inhaled glucocorticosteroids (antiasthmatics) vs. no treatment be used for the treatment of allergic rhinitis?				×	
Should cough suppressants vs. no treatment be used for the treatment of allergic rhinitis?				×	
Should expectorant medications (mucoactive agents) vs. no treatment be used for the treatment of allergic rhinitis?				×	
Should acetaminophen vs. no treatment be used for the treatment of allergic rhinitis?				×	
Should nonsteroidal anti-inflammatory drugs vs. no treatment be used for the treatment of allergic rhinitis?				×	
Should decongestant topical ointments vs. no treatment be used for the treatment of allergic rhinitis?				×	
Should specific sleep positions vs. other sleep positions be used for the management of allergic rhinitis?				×	
Should humidifiers vs. no intervention be used for the management of allergic rhinitis?				×	

(Continues)

TABLE 2 | (Continued)

Guideline question	Method for question generation				Prioritised
	ARIA 2010– 2016	Guideline panel/KOL	MASK-air data	Generation based on web search queries	Direct generation by LLM
Should nasal strips vs. no intervention be used for the management of allergic rhinitis?				×	
Should intranasal decongestants vs. oral decongestants be used for the treatment of allergic rhinitis?					×
Should intranasal ipratropium bromide vs. intranasal glucocorticosteroids be used for the treatment of allergic rhinitis?					×
Should psychological support or stress management techniques vs. no psychological intervention be used in patients with allergic rhinitis?					×
Should tailored advice on physical activity vs. no specific advice be used in patients with allergic rhinitis?					×
Should a systematic diagnostic screening for sleep disturbances, including sleep apnea, vs. no systematic screening be used in patients with allergic rhinitis?					×
Should telemedicine vs. in-person visits alone be used in patients with allergic rhinitis?					×

Abbreviations: ARIA, Allergic Rhinitis and its Impact on Asthma; KOL, key opinion leaders; LLM, large language models.

corresponding to the occurrence of any AE. The latter corresponded to the undesirable effect outcome with the second highest priority rating.

6 | Emphasising Diverse Populations

6.1 | Age Groups

For each guideline recommendation, we provide subgroup considerations for children. To gather evidence on the effectiveness and safety of AR treatments in children, we conducted systematic reviews solely assessing the paediatric population.

Between adolescents and adults, similar AR control levels were found using evidence from MASK-air [42]. On the other hand, there seem to be differences for individuals over 75 years of age [43], despite the lack of available evidence.

6.2 | Sex

Sex impacts health, with relevant effect modification and implications on the prevention, screening, diagnosis and treatment of allergic diseases [44]. This is particularly true during puberty and pregnancy [45–47]. However, since most studies do not provide evidence on the effect of interventions according to sex, more data are needed to incorporate sex-specific considerations in the guidelines.

6.3 | Patients With Multimorbid Asthma and/or Conjunctivitis

There is mounting evidence that AR alone and AR + asthma multimorbidity represent two distinct phenotypes [48] (Table S3). Such evidence—obtained from the MeDALL (Mechanisms of the Development of Allergy) study [49] and MASK-air data [50]—is grounded on insights into polysensitisation and multimorbidity, advances in mHealth for novel phenotype definition, confirmation in canonical epidemiologic studies and genomic findings. Therefore, subgroup considerations for patients with asthma were provided in the guidelines whenever justified.

Differences between AR alone and AR + conjunctivitis were identified with MASK-air data [51–54] and confirmed in canonical epidemiologic studies. These studies have shown that ocular symptoms (i) are more common in AR + asthma than in AR alone [54], (ii) are associated with the severity of nasal symptoms [55, 56] and (iii) are important to consider in severe asthma [55]. These data indicate that conjunctivitis should be considered as a separate disease in AR or AR + asthma and should be embedded in the multimorbid phenotype. However, more data are needed to incorporate this phenotype in guidelines.

7 | Obtaining Evidence for Evidence-to-Decision Frameworks

Recommendations were generated using EtDs [11, 12]. In its standard form, the EtD comprises 12 criteria, including

problem priority, desirable effects, undesirable effects, certainty of evidence [in desirable and undesirable effects], values and preferences, balance of effects, resources required, certainty of evidence of required resources, cost-effectiveness, equity, acceptability and feasibility [11, 12]. In addition, in ARIA-EAACI 2024–2025, a 13th criterion, Planetary Health, was considered [57]. The next subsections of this paper will discuss the evidence sources used to inform each EtD criterion. An example of an EtD (comparison between fixed combinations of INCS + intranasal antihistamines [INAH] vs. INCS) is available in the Appendix S1.

7.1 | Desirable and Undesirable Effects of Interventions

7.1.1 | Pairwise and Network Meta-Analyses of Randomised Controlled Trials

There was insufficient evidence regarding the comparative efficacy and safety of pharmacological treatments of AR before the beginning of this project. We therefore decided to conduct systematic reviews with pairwise or network meta-analyses to inform the comparative effectiveness of treatments [58]. Overall, for informing ARIA 2024–2025, we planned to conduct six systematic reviews of RCTs comparing the desirable and undesirable effects of oral and intranasal medications:

- i. Systematic review and pairwise meta-analysis comparing intranasal medications versus placebo in adults;
- ii. Systematic review and network meta-analysis comparing intranasal medications in adults;
- iii. Systematic review and network meta-analysis comparing intranasal medications in children;
- iv. Systematic review and pairwise meta-analysis comparing intranasal versus oral medications in adults;
- v. Systematic review and network meta-analysis comparing individual oral medications in adults and children;
- vi. Systematic review and network meta-analysis comparing classes of intranasal and oral medications in adults and children.

Three systematic reviews [59–61] and two protocols [62, 63] have been published. In line with the prioritised outcomes for ARIA-EAACI 2024–2025, these systematic reviews assessed the effects of each individual medication or medication class on nasal symptoms, ocular symptoms, rhinoconjunctivitis-related quality of life and AEs. Pairwise and network meta-analyses including both active comparisons and comparing medications with placebo provided information on untreated patients [59–61]. Network meta-analyses of active comparisons provided information on treated patients [59].

As for the main results of the published studies, both the pairwise and the network meta-analysis on intranasal medications have found that azelastine-fluticasone, fluticasone furoate and fluticasone propionate were the medications with the highest probability of resulting in moderate or large improvements in the assessed outcomes [59, 60]. Considering the differential

impact on ocular symptoms, fluticasone furoate or fluticasone propionate may be particularly indicated for patients with severe nasal symptoms but no or mild ocular symptoms, while azelastine-fluticasone may be an adequate option for patients with moderate to severe ocular symptoms.

In addition, the network meta-analysis on intranasal medications found that azelastine-fluticasone and fluticasone furoate were the medications with the highest probability of being the most efficacious, as they frequently resulted in clinically meaningful larger improvements when compared to other active treatments [59]. In children, evidence was scarcer than in adults. However, despite being efficacious and safe, the efficacy of AR treatments in children does not seem to be as high as in adults.

On the other hand, the meta-analysis comparing intranasal versus oral medications provided evidence that intranasal medications are more effective than oral ones (namely, oral antihistamines or oral leukotriene-receptor antagonists) [61].

Differences in the frequency of undesirable effects (AEs or serious AEs) were trivial for most comparisons between individual medications or between different medication classes [59, 61].

7.1.2 | Pharmacovigilance Data

Considering limitations of RCTs in registering AEs (e.g., limited number of participants, short period of follow-up and restricted eligibility criteria) [64, 65], we analysed pharmacovigilance data to provide further information on the undesirable effects of interventions. We queried the VigiBase database on reported AEs to individual medications of different classes used for the treatment of AR. VigiBase, the World Health Organization global database of AE reports for medicines and vaccines, is the largest pharmacovigilance database [66]. The analysis of its data allowed us to identify the 15 most commonly reported AEs for each intervention being compared in each question. For each AE, we calculated the respective reporting odds ratio to identify signals of disproportionate reporting [67]. For example, in the comparison between INCS and INAH, we found that cataract and glaucoma were proportionally more commonly reported with INCS than with INAH.

7.2 | Values

Healthcare interventions typically result in benefits and harms. Values concern the relative importance patients place on specific benefits and harms [38, 68]. We performed a systematic review to provide a comprehensive overview of the values of patients with AR [38]. We observed that (i) patients generally value the efficacy of interventions more than their AEs, and (ii) patients consider nasal symptoms (particularly nasal obstruction) as those with the highest impact. One of the included studies of that systematic review was based on direct patient data from MASK-air, involving the computation of utilities in different European countries according to the level of rhinitis control and the presence of comorbid asthma [69]. The results of the study were coherent with a questionnaire sent to ARIA members [70].

7.3 | Resources Required and Cost-Effectiveness

7.3.1 | Global Survey on Medication Costs

The cost of medications is highly variable across countries. Moreover, medication costs change with time, particularly when generic drugs are made available or when medications become over-the-counter. Therefore, to gather information on this criterion, we conducted a survey asking ARIA experts about the availability and lowest costs of individual rhinitis medications in their countries (Figure 3).

7.3.2 | MASK-air Data on Indirect Costs Associated With Rhinitis Control

We used data from MASK-air to quantify indirect costs resulting from productivity losses associated with worse rhinitis control. MASK-air includes WPAI:AS, a validated questionnaire that measures the impact of allergies on absenteeism and presenteeism [32]. Based on these data, we estimated the underlying indirect costs in 40 countries according to the level of rhinitis control [71]. Overall, median work impairment (including both absenteeism and presenteeism) stood at 4.6% (P25-P75 = 0.8%–15.1%) for well-controlled weeks, ascending to 27.7% (P25-P75 = 12.4%–46.0%) and 60.7% (P25-P75 = 38.5%–80.2%) for partially and poorly controlled weeks, respectively. For example, in Germany, this translates into median weekly monetary losses of 42.3 US dollars (US\$) adjusted for purchasing power parity (PPPs) for well-controlled weeks, 259.7 US\$ PPPs for partially controlled weeks, and 554.2 US\$ PPPs for poorly controlled weeks (difference of 511.9 US\$ per week from poor to good control). For other Western European countries, the difference ranges from 200 to 695 US\$ [71].

7.3.3 | “Cost-Utility” Assessments Based on Health Technology Assessment Reports and MASK-air Data

There is a scarcity of recent cost-effectiveness and cost-utility studies on AR treatments. We therefore considered cost data from the survey to ARIA experts and either (i) utilities data estimated from a health technology assessment report [72] or (ii) MASK-air data on levels of the VAS of EQ-5D associated with each intervention. The use of VAS EQ-5D data reported in MASK-air was justified since (i) the identified health technology assessment report did not assess all medication classes and (ii) there is a lack of available studies estimating utilities associated with different treatments for AR.

7.4 | Equity

7.4.1 | Global Survey on the Availability of Medications

To assess across-country differences in the availability of medications for AR, we conducted a survey asking ARIA experts about the availability of individual rhinitis medications in their countries (Figure 3).




FIGURE 3 | Countries with valid answers to the survey sent to ARIA experts on the availability and costs of allergic rhinitis medication. The answers provided up until September 2024 were used to inform the ARIA-EAACI 2024–2025 guidelines.

7.4.2 | World Health Organization List of Essential Medicines

We assessed the medications for AR included in the World Health Organization List of Essential Medicines [73]. Only intranasal budesonide, oral loratadine and intranasal xylometazoline are on this list. According to information from local experts, medications available in this list are provided free of charge in some low-income countries.

7.5 | Acceptability and Feasibility

7.5.1 | Assessment of Patient Satisfaction With Treatment in MASK-air Data

Patients' satisfaction with their treatment is a patient-reported outcome measure distinct from quality of life and symptoms [74]. In its daily monitoring questionnaire, the MASK-air app includes a question assessing (through a 0–100 VAS) how satisfied patients are with their rhinitis treatment on that day. We have analysed VAS satisfaction data for each treatment class or individual medication in single medication and in co-medication. In addition, we have built multivariable regression models to compare different medications on VAS satisfaction levels.

7.5.2 | Assessment of Co-Medication Use in the MASK-air Data

We assessed MASK-air data to compare different individual medications and medication classes on the frequency of their use in co-medication. MASK-air studies have consistently found

that when patients use co-medication, they tend to report more severe rhinitis symptoms [18, 75]. This finding may be explained by the fact that patients tend to increase their medication use when feeling less well controlled.

7.5.3 | Assessment of Patients' Adherence to Treatment in the MASK-air Data

RCTs tend to assess strategies involving the daily use of medications and, henceforth, require a high adherence to treatment (usually 70%). However, in MASK-air, two studies have addressed adherence to rhinitis treatment and have found that most AR patients are not adherent [76, 77]. In particular, most patients use treatment on an as-needed basis when feeling symptomatic [18]. Therefore, to inform about the acceptability of interventions, we analysed MASK-air data to compare the adherence to different rhinitis medication classes.

7.5.4 | Speed of Onset of Action of Medications: Rapid Evidence Review

We conducted a rapid evidence review to assess the speed of onset of action of AR medications based on ARIA 2019 and other sources [9]. We took into account the guideline published by the European Medicines Agency on the “clinical development of medicinal products for the treatment of allergic rhinoconjunctivitis” [78], and considered the onset of action of medications. Evidence on the onset of action of medications for AR may originate from three study types: standard phase III double-blind RCTs, park setting studies (natural exposure) and allergen exposure chamber studies (artificial exposure). In this rapid review, we found that medications

containing INAH act within 5–30mins, whereas most INCS need hours to be effective.

7.6 | Planetary Health

The concept of Planetary Health emphasises the inherent connection between the health of humans and the health of the planet, on which human health depends [79]. There is increasing understanding that clinical decision guidelines need to incorporate a planetary health dimension [80]. The consideration of Planetary Health in ARIA started with the participation in an EU High-Level meeting (Finnish Presidency of the European Union and DG Research) on Planetary Health [81–83].

ARIA-EAACI 2024–2025 considered aspects of Planetary Health in the formulation of guideline recommendations by having Planetary Health criteria in the EtD [57, 80]. For example, for comparisons between intranasal versus oral treatments, aspects such as the global warming potential of the different packaging types and manufacturing places were considered for available evidence.

8 | Issuing Judgements and Actionable, Patient-Centred Recommendations for Each Guideline Question

In the EtDs, evidence is provided for each criterion, so that the guideline panel can make a judgement. Based on the judgements provided for all criteria, the panel is then able to issue a recommendation for the respective guideline question.

The ARIA-EAACI 2024–2025 guideline panel held regular online meetings to make judgements on the EtD criteria and formulate guideline recommendations. In each case, consensus among members of the panel without CoI was sought. If consensus was not reached, a formal voting process was set (using the anonymous voting feature of GRADEpro), although restricted to guideline panel members without CoI.

We drafted recommendations as suggested by the GRADE working group. GRADE recommendations are characterised by (i) a directionality (i.e., whether there is favouring of the intervention, of the comparison, or of neither the intervention nor the comparison) and (ii) a strength (i.e., whether the recommendation is strong or conditional). Strong recommendations posit




FIGURE 4 | Example of an electronic clinical decision support system (ARIA 2016) (example for patients with well-controlled allergic rhinitis). Ti = Class of current treatment (in case of polypharmacy, Ti = maximum class); Tj, Tk and Tl = Medications added to Ti, order of class $1 < k < j < i$. T1 = antihistamine (oral, intranasal and eye drops), leukotriene-receptor antagonist, chromone (intranasal and eye drops); T2 = intranasal corticosteroid (INCS); T3 = INCS + azelastine; T4 = add short course of oral corticosteroids; T5 = consider referral and allergen-specific immunotherapy.

that the intervention should be applied to most individuals. Conditional recommendations indicate that the recommendation may not be applicable to relevant subsets of patients and that variability in the clinical practice may be appropriate [84, 85]. In addition, recommendations should inform on the underlying certainty of evidence (quality of the whole body of evidence, considering altogether desirable and undesirable effects).

After drafting guideline recommendations, the guideline panel proposes subgroup considerations for children and, if justified, patients with asthma. If necessary, the guideline panel also (i) suggests implementation considerations (e.g., in aspects related to the application of recommendations in low- and middle-income countries), (ii) discusses aspects related to monitoring and evaluation and/or (iii) highlights related research priorities.

9 | From Recommendations to Action: Building Digitalised Decision Algorithms

In ARIA 2010–2016, a simple algorithm was proposed in treated and untreated patients. In ARIA-EAACI 2024–2025, we will propose more management algorithms, considering a diverse set of scenarios in terms of patients (e.g., controlled and treated, uncontrolled and treated or uncontrolled and untreated), settings (e.g., primary care or specialised allergy care) or regions (e.g., low- and middle-income countries or high-income countries; the survey we conducted among ARIA experts will support such tailoring, as it informs on the availability and price of medication in different countries). The scenarios will be jointly proposed by the guideline panel group, the review group and ARIA Junior members. The corresponding algorithms will be built based on the formulated recommendations.

In ARIA 2010–2016, an algorithm was developed [14]. Moreover, an electronic clinical decision system provided evidence-based treatment recommendations. The app interfaces used to collect the necessary information had been described, and the management algorithms had been programmed into the MASK-air app using Titanium Appcelerator for iOS tablets [86] (Figure 4 and Figure S1). However, since MASK-air was not at that time an MDR Class IIa, the digitalised algorithms were not implemented. In ARIA 2024–2025, the developed management algorithms will be digitalised and implemented in MASK-air.

10 | Conclusions

ARIA-EAACI 2024–2025 will provide major advances for the management of patients with AR. Building on several different sources of evidence, ARIA-EAACI 2024–2025 will adopt a patient-centred perspective and promote shared decision-making. The formulation of recommendations will be fully based on the GRADE working group methodology. Nevertheless, the development of ARIA-EAACI 2024–2025 will contribute per se to methodological innovations for the guideline development process in aspects such as the use of AI to support the development of questions, or the incorporation of evidence from multiple complementary sources (Table 1). As its end product, ARIA-EAACI 2024–2025 aims not only at producing a set of recommendations but also at developing a set of digitalised

management algorithms, which will be essential in promoting a continuative and long-standing integration of these guidelines into daily care.

Affiliations

¹Institute of Allergology, Charité – Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany | ²Fraunhofer Institute for Translational Medicine and Pharmacology ITMP, Immunology and Allergology, Berlin, Germany | ³ARIA (Allergic Rhinitis and Its Impact on Asthma), Montpellier, France | ⁴MEDCIDS—Department of Community Medicine, Information and Health Decision Sciences, Faculty of Medicine, University of Porto, Porto, Portugal | ⁵CINTESIS@RISE—Health Research Network, Faculty of Medicine, University of Porto, Porto, Portugal | ⁶Department of Health Research Methods, Evidence, and Impact & Department of Medicine, McMaster University, Hamilton, Ontario, Canada | ⁷Department of Biomedical Science, Humanitas University, Pieve Emanuele, Milan, Italy | ⁸Division of Allergy, Immunology, and Transplantation (DAIT), National Institute of Allergy and Infectious Diseases, NIH, Bethesda, MD, Maryland, USA | ⁹Department of Prevention of Environmental Hazards, Allergology and Immunology, Medical University of Warsaw, Warsaw, Poland | ¹⁰Institute of Clinical Medicine and Institute of Health Sciences, Medical Faculty of Vilnius University, Vilnius, Lithuania | ¹¹Clinic of Asthma, Allergy, and Chronic Lung Diseases, Asthma & Allergy Department, Vilnius, Lithuania | ¹²International Primary Care Respiratory Group IPCRG, Edinburgh, UK | ¹³Medical Consulting Czarlewski, Levallois, France | ¹⁴Allergy Unit, Málaga Regional University Hospital of Málaga, Malaga University, ARADyAL, Malaga, Spain | ¹⁵National Heart and Lung Institute (NHLI), Imperial College, and NIHR Imperial Biomedical Research Centre, London, UK | ¹⁶Allergy Center, CUF Descobertas Hospital, Lisbon, Portugal | ¹⁷Asthma & Allergy Unit, IRCCS Humanitas Research Hospital, Rozzano, Milan, Italy | ¹⁸Department of Allergy, Hospital La Paz Institute for Health Research (IdiPAZ), Madrid, Spain | ¹⁹Section of Allergy and Immunology, Saint Louis University School of Medicine, Saint Louis, Missouri, USA | ²⁰“Respiriamo Insieme” Association, Asthma & Allergy Center, Padova, Italy | ²¹Department of Otolaryngology, Head & Neck Surgery, Universitätsmedizin Mainz, Mainz, Germany | ²²Center for Rhinology and Allergology, Allergology & Rhinology Department, Wiesbaden, Germany | ²³Department of Political Science, University of Zürich, Zürich, Switzerland | ²⁴RISE-Health, Department of Medical Sciences, Faculty of Health Sciences, University of Beira Interior, Covilhã, Portugal | ²⁵Finnish Meteorological Institute (FMI), Helsinki, Finland | ²⁶Allergy Department, 2nd Pediatric Clinic, National and Kapodistrian University of Athens, Athens, Greece | ²⁷PaCeIT—Patient Centered Innovation and Technologies, Center for Health Technology and Services Research (CINTESIS), Faculty of Medicine, University of Porto, Porto, Portugal | ²⁸Allergy Unit, Instituto and Hospital CUF, Porto, Portugal | ²⁹Barcelona Institute for Global Health (ISGlobal), Barcelona, Spain | ³⁰Universitat Pompeu Fabra (UPF), Barcelona, Spain | ³¹CIBER Epidemiología y Salud Pública (CIBERESP), Barcelona, Spain | ³²Department of Allergy, Skin and Allergy Hospital, Inflammation Center, Helsinki University Hospital and University of Helsinki, Helsinki, Finland | ³³Department of Otorhinolaryngology, University of Eastern Finland and the North Savo Wellbeing Services County, Kuopio, Finland | ³⁴Allergy and Clinical Immunology, University of Bari Medical School, Bari, Italy | ³⁵Institute of Sciences of Food Production, National Research Council (ISPA-CNR), Bari, Italy | ³⁶School of Medicine, Universidad de los Andes, Bogotá, DC, Colombia | ³⁷Pulmonology Service, Internal Medicine Section, Fundación Santa Fe de Bogotá, University Hospital, Bogotá, DC, Colombia | ³⁸Fundação ProAR and (Faculdade de Medicina da), Universidade Federal da Bahia, Salvador, Bahia, Brazil | ³⁹Casa di Cura Villa Montallegro, Allergology Department, Genova, Italy | ⁴⁰Department of Pulmonary Diseases and Institute of

Pulmonology and Tuberculosis, Istanbul University-Cerrahpaşa, Cerrahpaşa Faculty of Medicine, Istanbul, Turkey | ⁴¹Department of Health Sciences, University of Florence, Florence, Italy | ⁴²Allergy Unit, Meyer Children's Hospital IRCCS, Florence, Italy | ⁴³Department of Immunology and Allergology, Lithuanian University of Health Sciences, Kaunas, Lithuania | ⁴⁴Department of Pediatrics and Adolescent Medicine, University of Turku, and Turku University Hospital, Turku, Finland | ⁴⁵Institute of Clinical Immunology and Medical Genetics, Department of Paediatrics and Adolescent Medicine, Department of Pulmonology and Phthisiology, Jessenius Faculty of Medicine, Martin, Slovakia | ⁴⁶Comenius University, Bratislava, Slovakia | ⁴⁷University Teaching Hospital Martin, Martin, Slovakia | ⁴⁸Division of Internal Medicine, Asthma and Allergy, Barlicki University Hospital, Medical University of Lodz, Lodz, Poland | ⁴⁹Institute of Clinical Medicine, Clinic of Chest Diseases and Allergology, Faculty of Medicine, Vilnius University, Vilnius, Lithuania | ⁵⁰Institute of Biomedical Sciences, Department of Pathology, Faculty of Medicine, Vilnius University, Vilnius, Lithuania | ⁵¹Center of Excellence in Asthma and Allergy, Médica Sur Clinical Foundation and Hospital, México City, Mexico | ⁵²Allergy & Immunology Centre, Pantai Hospital Kuala Lumpur, Kuala Lumpur, Malaysia | ⁵³National Center for Research in Chronic Respiratory Diseases, Collaborating With WHO—EMRO Tishreen University School of Medicine, Latakia, Syria | ⁵⁴Al-Sham Private University, Damascus, Syria | ⁵⁵Japan Antituberculosis Association (JATA), Fukuji Hospital, Tokyo, Japan | ⁵⁶Department of Respiratory Medicine, JSS Medical College, JSSAHER, Mysore, Karnataka, India | ⁵⁷Departement of Biological Sciences and Pathophysiology, University of Veterinary Medicine, Vienna, Austria | ⁵⁸Section of Rhinology and Allergy, Department of Otorhinolaryngology, Head and Neck Surgery, University Hospital Marburg, Philipps-Universität Marburg, Marburg, Germany | ⁵⁹Allergy and Clinical Immunology Department, Hospitais da Universidade de Coimbra, Unidade Local de Saúde de Coimbra, Coimbra, Portugal | ⁶⁰Center for Innovative Biomedicine and Biotechnology (CIBB), Faculty of Medicine, University of Coimbra, Coimbra, Portugal | ⁶¹Institute of Immunology, Faculty of Medicine, University of Coimbra, Coimbra, Portugal | ⁶²UBIAir—Clinical & Experimental Lung Centre and CICS-UBI Health Sciences Research Centre, University of Beira Interior, Covilhã, Portugal | ⁶³Pneumologie, AP-HP Centre Université de Paris Cité, Hôpital Cochin, Paris, France | ⁶⁴UMR 1016, Institut Cochin, Paris, France | ⁶⁵Inserm, Equipe d'Epidémiologie Respiratoire Intégrative, CESP, Villejuif, France | ⁶⁶Department of Immunoallergology, Cova da Beira University Hospital Centre, Covilhã, Portugal | ⁶⁷UBIAir—Clinical & Experimental Lung Centre, University of Beira Interior, Covilhã, Portugal | ⁶⁸Department of Respiratory Medicine, Copenhagen University Hospital-Hvidovre, Copenhagen, Denmark | ⁶⁹Institute of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark | ⁷⁰Pediatrics, Allergy & Immunology, Latin American Society of Allergy, Asthma & Immunology (SLAAI), Montevideo, Uruguay | ⁷¹Pulmonary Environmental Epidemiology Unit, CNR Institute of Clinical Physiology, Pisa, Italy | ⁷²Department of Otolaryngology, Head and Neck Surgery, Beijing TongRen Hospital and Beijing Institute of Otolaryngology, Beijing, China | ⁷³ISGlobal, Barcelona Institute for Global Health, Barcelona, Spain | ⁷⁴Skin and Allergy Hospital, Helsinki University Hospital and University of Helsinki, Helsinki, Finland | ⁷⁵Universidad Espíritu Santo, Samborondón, Ecuador | ⁷⁶Respiralab Research Group, Guayaquil, Guayas, Ecuador | ⁷⁷Servicio de Alergia e Immunologia, Clinica Santa Isabel, Buenos Aires, Argentina | ⁷⁸Global NCD Platform, Geneva, Switzerland | ⁷⁹Department of Pulmonary Diseases, Celal Bayar University, Faculty of Medicine, Manisa, Turkey | ⁸⁰Department of Otorhinolaryngology—Head and Neck Surgery, School of Medical Sciences, Universiti Sains Malaysia, Kubang Kerian, Kelantan, Malaysia | ⁸¹Microbiology Department, College of Medicine, Kuwait University, Kuwait City, Kuwait | ⁸²Adult Allergy and Immunology Division—Hamad Medical Corporation, Doha, Qatar | ⁸³Department of Cardiovascular and Respiratory Sciences, Porto Health School, Polytechnic Institute of Porto, Porto, Portugal | ⁸⁴Department of Women's and Children's Health, Paediatric Research, Uppsala

University, Uppsala, Sweden | ⁸⁵Department of Internal Medicine, University of Medicine, Tirana, Albania | ⁸⁶Division of Immunology, Allergy and Rheumatology, Department of Medicine, University of Cincinnati College of Medicine, Cincinnati, Ohio, USA | ⁸⁷Department of Pediatrics, Medical College of Georgia at Augusta University, Augusta, Georgia, USA | ⁸⁸Department of Internal Medicine (DIMI), University of Genoa, Genoa, Italy | ⁸⁹Respiratory and Allergy Clinic, IRCCS - Policlinico San Martino, Genoa, Italy | ⁹⁰Department of Pediatrics, Medical School, Federal University of Minas Gerais, Belo Horizonte, Brazil | ⁹¹CHRC, LA-REAL, NOVA Medical School, NMS, Universidade NOVA de Lisboa, Lisboa, Portugal | ⁹²Serviço de Imunoalergologia, Hospital de Dona Estefânia, Lisbon, Portugal | ⁹³Division of Allergy/Immunology, University of South Florida, Tampa, Florida, USA | ⁹⁴SOS Allergology and Clinical Immunology, USL Toscana Centro, Prato, Italy | ⁹⁵Allergy, Bambino Gesù Children's Hospital, Istituto di Ricovero e Cura a Carattere Scientifico (IRCCS), Rome, Italy | ⁹⁶Department of Internal Medicine 'A. Murri' and Unit of Geriatric Immunoallergology, University of Bari Medical School, Bari, Italy | ⁹⁷Faculty of Public Health, Sofia Medical University, Sofia, Bulgaria | ⁹⁸Latvian Association of Allergists, University Children Hospital, Riga, Latvia | ⁹⁹Life and Health Sciences Research Institute (ICVS), School of Medicine, University of Minho, Braga, Portugal | ¹⁰⁰CINTESIS@RISE, Biochemistry Lab, Faculty of Pharmacy and Competence Center on Active and Healthy Ageing, University of Porto, Porto, Portugal | ¹⁰¹Department of Medical Sciences and Public Health and Unit of Allergy and Clinical Immunology, University Hospital "Duisio Casula", University of Cagliari, Cagliari, Italy | ¹⁰²VIM Suresnes, UMR 0892, Pôle des Maladies des Voies Respiratoires, Hôpital Foch, Université Paris-Saclay, Suresnes, France | ¹⁰³Medical Faculty Skopje, University Clinic of Pulmonology and Allergy, Skopje, Republic of Macedonia | ¹⁰⁴Pediatric Allergy, Immunology and Rheumatology Unit, Children's Hospital, Ain Shams University, Cairo, Egypt | ¹⁰⁵Department of Otorhinolaryngology, Chiba University, Chiba, Japan | ¹⁰⁶Department of Internal Medicine and Infectious Diseases, St Joseph University, Hotel Dieu de France Hospital, Beirut, Lebanon | ¹⁰⁷Department of Allergology and Clinical Immunology, Kazakhstan Association of Allergology and Clinical Immunology, Kazakh National Medical University, Almaty, Kazakhstan | ¹⁰⁸Institute of Clinical Medicine, Children's Clinic, Tartu University, Tartu, Estonia | ¹⁰⁹Poltava State Medical University, Immunology & Allergology Department, Poltava, Ukraine | ¹¹⁰Department of Respiratory Medicine, National Institute of Diseases of the Chest and Hospital, Dhaka, Bangladesh | ¹¹¹Department of Otorhinolaryngology, Head and Neck Surgery, Semmelweis University, Budapest, Hungary | ¹¹²Department of Clinical Science and Education, Södersjukhuset, Karolinska Institutet, Stockholm, Sweden | ¹¹³Sach's Children and Youth Hospital, Södersjukhuset, Stockholm, Sweden | ¹¹⁴Department of Pediatric Respiratory Diseases and Allergology, Medical University of Warsaw, Warsaw, Poland | ¹¹⁵Department of Clinical and Laboratory Immunology, Allergology and Medical Genetics, Bogomolets National Medical University, Kyiv, Ukraine | ¹¹⁶Institute of Translational Pharmacology (IFT)-National Research Council (CNR), Palermo, Italy | ¹¹⁷Croatian Pulmonary Society, Clinical Center for Pulmonary Diseases, Zagreb, Croatia | ¹¹⁸Asthma, COPD Outpatient Care Unit, University Medical Center, Hô-Chi-Minh, Vietnam | ¹¹⁹Allergy Unit "D Kalogeromitros", 2nd Department of Dermatology and Venereology, National & Kapodistrian University of Athens, "Attikon" University Hospital, Athens, Greece | ¹²⁰Clinic for Pulmonary Diseases, Clinical Center of Serbia, Faculty of Medicine, University of Belgrade, Serbian Association for Asthma and COPD, Belgrade, Serbia | ¹²¹Division of Respiratory Disease and Allergy, Department of Internal Medicine, Dankook University College of Medicine, Cheonan, Republic of Korea | ¹²²Department of Medicine, Faculty of Medicine and Surgery, University of Malta, MSD, Msida, Malta | ¹²³EPIUnit-Institute of Public Health, University of Porto, and Laboratory for Integrative and Translational Research in Population Health (ITR), Porto, Portugal | ¹²⁴Serviço de Imunoalergologia, Centro Hospitalar Universitário São João, Porto, Portugal | ¹²⁵Basic and Clinical Immunology Unit, Department of Pathology, Faculty of Medicine,

University of Porto, Porto, Portugal | ¹²⁶Rhinology Unit & Smell Clinic, Department of Otorhinolaryngology, Hospital Clínic Barcelona, Universitat de Barcelona; FRCB-IDIBAPS; CIBERES, Barcelona, Spain | ¹²⁷Université Paris-Saclay, UVSQ, Univ. Paris-Sud, Villejuif, France | ¹²⁸Inserm, Équipe d'Epidémiologie Respiratoire Intégrative, Villejuif, France | ¹²⁹Imperial College Healthcare NHS Trust, London, UK | ¹³⁰University of Liverpool, Liverpool, UK | ¹³¹Center of Allergy, Immunology and Respiratory Diseases, Santa Fe, Argentina | ¹³²Department of Allergology, Medical University of Gdansk, Gdansk, Poland | ¹³³ENT Department, University Hospital of Kinshasa, Kinshasa, Democratic Republic of Congo | ¹³⁴Allergy, Asthma and Clinical Immunology, Alfred Health, Department of Immunology, Central Clinical School, Monash University, Melbourne, Victoria, Australia | ¹³⁵Swiss Institute of Allergy and Asthma Research (SIAF), University of Zurich, Davos, Switzerland | ¹³⁶Chiba Rosai Hospital, ENT Department, Chiba, Japan | ¹³⁷Chiba University Hospital, Department of Otolaryngology, Head and Neck Surgery, Chiba, Japan | ¹³⁸Department of Otorhinolaryngology, Charité – Universitätsmedizin Berlin, Berlin, Germany | ¹³⁹Department of Biochemistry and Molecular Biology, School of Chemistry, Complutense University of Madrid, Madrid, Spain | ¹⁴⁰Department of Immunology and Allergology, Faculty of Medicine in Pilsen, Charles University, Prague, Czech Republic | ¹⁴¹Division of Allergy and Clinical Immunology, Department of Medicine, “Santa Maria della Speranza” Hospital, Battipaglia, Salerno, Italy | ¹⁴²Agency of Health ASL, Division of Allergy and Clinical Immunology, Department of Medicine, Salerno, Italy | ¹⁴³Postgraduate Programme in Allergy and Clinical Immunology, University of Naples Federico II, Naples, Italy | ¹⁴⁴Department of Pediatrics, Nippon Medical School, Tokyo, Japan | ¹⁴⁵Medical School, University of Cyprus, Nicosia, Cyprus | ¹⁴⁶Clinic of Occupational Diseases, University Hospital Sveti Ivan Rilski, Sofia, Bulgaria | ¹⁴⁷IRCCS Humanitas Research Center, Personalized Medicine Asthma & Allergy, Rozzano, Milan, Italy | ¹⁴⁸Department of Pulmonology and Allergology, Klaipeda National Hospital, Klaipeda, Lithuania | ¹⁴⁹Vilnius University Medical Faculty, Department of Political Behaviour and Institutions, Vilnius, Lithuania | ¹⁵⁰Division of Adult and Pediatric Allergy and Immunology, University of the Philippines - Philippines General Hospital, Manila, Philippines | ¹⁵¹Department of Allergy, Clinics Hospital, National University, San Lorenzo, Paraguay | ¹⁵²Faculty of Medicine, University of Southampton, Southampton, UK | ¹⁵³The David Hide Asthma and Allergy Centre, St Mary's Hospital, Isle of Wight, UK | ¹⁵⁴NIHR Southampton Biomedical Research Centre, University Hospital Southampton NHS Foundation Trust, Southampton, UK | ¹⁵⁵LEADER Research Inc, Hamilton, Ontario, Canada | ¹⁵⁶Division of Allergy, Asthma and Clinical Immunology, Emek Medical Center, Afula, Israel | ¹⁵⁷Rappaport Faculty of Medicine, Technion-Israel Institute of Technology, Haifa, Israel | ¹⁵⁸ARIA, Asthma, Rhinitis, Immunology & Allergy Department, Athens, Greece | ¹⁵⁹Allergy Service, Fundacion Jimenez Diaz, Autonoma University of Madrid, CIBERES-ISCIII, Madrid, Spain | ¹⁶⁰PROMISE Department, University of Palermo, Palermo, Italy | ¹⁶¹Allergy & Asthma, Medical Director, CLINICA SISUL, FACAIA, SPAIA, Asuncion, Paraguay | ¹⁶²Division of Allergy, Clinical Immunology and Rheumatology, Department of Pediatrics, Federal University of São Paulo, São Paulo, Brazil | ¹⁶³Division of Respiratory Medicine, Department of Pediatrics, Hospital Nacional de Niños, Universidad de Costa Rica, San Jose, Costa Rica | ¹⁶⁴Department of Respiratory Medicine and Tuberculosis, University Hospital, Brno, Czech Republic | ¹⁶⁵Department of Otolaryngology, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok, Thailand | ¹⁶⁶Imunoalergologia, Centro Hospitalar Universitário de Coimbra, Faculty of Medicine, University of Coimbra, Coimbra, Portugal | ¹⁶⁷International Primary Care Respiratory Group IPCRG, Aberdeen, UK | ¹⁶⁸Health Planning Unit, Department of Social Medicine, Faculty of Medicine, University of Crete, Heraklion, Greece | ¹⁶⁹Department of Medicine, Federal University of Pampa, Uruguaiana, Brazil | ¹⁷⁰Department of Lung Diseases and Clinical Immunology, University of Turku, Turku, Finland | ¹⁷¹Research on Healthcare Performance (RESHAPE), INSERM U1290, Pneumologie et Médecine Respiratoire, Groupement Hospitalier Nord, Hôpital de la

Croix Rousse, Lyon, France | ¹⁷²FILHA, Finnish Lung Health Association, Helsinki, Finland | ¹⁷³Department of Clinical Medicine, Pulmonary Diseases and Clinical Allergology, University of Turku, Turku, Finland | ¹⁷⁴Nova Southeastern University, College of Allopathic Medicine, Fort Lauderdale, Florida, USA | ¹⁷⁵Department of Otolaryngology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Republic of Singapore | ¹⁷⁶Division of Allergy and Immunology, Department of Dermatology, Allergy and Venerology, Charité – Universitätsmedizin Berlin, Berlin, Germany | ¹⁷⁷The Allergy and Asthma Institute, Allergy & Asthma Department, Islamabad, Pakistan | ¹⁷⁸University Clinic of Respiratory and Allergic Diseases, Pulmonary & Allergy Department, Golnik, Slovenia | ¹⁷⁹Faculty of Medicine, University of Ljubljana, Ljubljana, Slovenia | ¹⁸⁰Pulmonary Department, Rashid Hospital, DUBAI Health, Dubai, UAE | ¹⁸¹Research Institute of Medical and Health Sciences, University of Sharjah, Sharjah, UAE | ¹⁸²Department of Allergy and Immunology, Hospital Quironsalud Bizkaia, Bilbao, Spain | ¹⁸³Oasi Research Institute-IRCCS, Troina, Italy | ¹⁸⁴BIOS S.p.A. Società Benefit, Rome, Italy | ¹⁸⁵Department of Allergy and Immunology, Para State University Center - CESUPA, Belém, Brazil | ¹⁸⁶Department for Otorhinolaryngology, Head and Neck Surgery, University of Tübingen, Tübingen, Germany | ¹⁸⁷Respiratory Medicine, University of Modena & Reggio Emilia, Azienda Ospedaliera-Universitaria di Modena, Modena, Italy | ¹⁸⁸Laval University, Quebec City, Quebec, Canada | ¹⁸⁹Department of Pediatric Allergology, Armand Trousseau University Hospital, Sorbonne University, AP-HP, Paris, France | ¹⁹⁰French National Reference Center for Angioedema (CREAK), Saint-Antoine University Hospital, Paris, France | ¹⁹¹CRESS, Inserm, INRAE, HERA Team, Paris Cité University, Paris, France | ¹⁹²Department of Medical Sciences, University of Turin, Turin, Italy | ¹⁹³Allergy and Clinical Immunology Unit, Mauriziano Hospital, Torino, Italy | ¹⁹⁴Department of Pulmonary Medicine, Mainz University Hospital, Mainz, Germany | ¹⁹⁵Pediatrics Department, Faculty of Medicine, Universidad Austral de Chile, Valdivia, Chile | ¹⁹⁶Allergy Section, Department of Internal Medicine, Hospital Vall D'hebron, Barcelona, Spain | ¹⁹⁷ARADyAL Research Network, Barcelona, Spain | ¹⁹⁸Pneumology Department, Hospital Universitari Dexeus, Barcelona, Spain | ¹⁹⁹CIBER of Respiratory Diseases, Group of Rhinitis, Rhinosinusitis and Nasal Polyps, Area of Asthma, Barcelona, Spain | ²⁰⁰Clinique des Bronches, Allergie et Sommeil, Hôpital Nord, Marseille, France | ²⁰¹National eHealth Living Lab, Department of Public Health and Primary Care, Leiden University Medical Center, Leiden, the Netherlands | ²⁰²Department of Allergology, Medical University of Gdańsk, Gdańsk, Poland | ²⁰³Department of Allergology and Department of Otorhinolaryngology, the First Affiliated Hospital, Nanjing Medical University, Nanjing, China | ²⁰⁴Allergist, David Tvildiani Medical University, Tbilisi, Georgia | ²⁰⁵Associate Professor of Pediatrics, Division of Allergy and Immunology, Federal University of Parana, Curitiba, Brazil | ²⁰⁶Paediatric Allergy Clinic, Department of Dermatology, Amersham Hospital - NHS Hospital Trust, Amersham, UK | ²⁰⁷Department of Health Research Methods, Evidence & Impact, McMaster University, Hamilton, Ontario, Canada | ²⁰⁸Evidence in Allergy Group, McMaster University and the Research Institute of St. Joe's Hamilton, Hamilton, Ontario, Canada | ²⁰⁹Medical Faculty, ENT Department, Eskisehir Osmangazi University, Eskisehir, Turkey | ²¹⁰Scientific & Medical Department, Lofarma S.p.A, Milan, Italy | ²¹¹Woolcock Institute of Medical Research, Sydney, Australia | ²¹²Allergy Service, University Hospital Professor Polydoro Ernani de São Thiago (HU-UFSC/EBSEH), Florianópolis, Brazil | ²¹³Department of Internal Medicine, Federal University of Santa Catarina (UFSC), Florianópolis, Brazil | ²¹⁴Division of Respiratory and Allergic Diseases, High Specialty Hospital 'a Cardarelli', and Respiratory Allergy School of Specialization in Respiratory Diseases, Federico II University of Naples, Naples, Italy | ²¹⁵Centre for Immunology and Infection Control, School of Biomedical Sciences, Faculty of Health, Queensland University of Technology, Brisbane, Australia | ²¹⁶Office of Research, Metro North Hospital and Health Service, Brisbane, Australia | ²¹⁷Department of Medical and Surgical Sciences, School of Allergology and Clinical Immunology, University of Foggia, Foggia, Italy | ²¹⁸Dr Agostinho Neto University Hospital,

Epidemiology Department, Praia, Cabo Verde | ²¹⁹Immunology, Cabo Verde University, Faculty of Medicine, Praia, Cape Verde | ²²⁰Department of Otorhinolaryngology, Head and Neck Surgery, University of Crete, School of Medicine, Heraklion, Crete, Greece | ²²¹Allergist, Private Practice, Mexico City, Mexico | ²²²Department of Internal Diseases, Allergy and Clinical Immunology, Medical University of Silesia in Katowice, Katowice, Poland | ²²³Department of Pulmonary Medicine, Division of Immunology, Allergy and Asthma, Laboratory of Occupational and Environmental Respiratory Diseases, Faculty of Medicine, Ege University, EgeSAM (Ege University Translational Pulmonary Research, Center), Bornova, Türkiye | ²²⁴Faculty of Health Sciences, Catholic University of Salta, Salta, Argentina | ²²⁵Allergy and Clinical Immunology Centro Regional, Hospital Universitario, Universidad Autónoma de Nuevo Leon, Monterrey, Mexico | ²²⁶Center of Allergy and Immunology, and Georgian Academy of Allergy, Asthma, and Clinical Immunology, Tbilisi, Georgia | ²²⁷Department of Allergy and Clinical Immunology, Air Force General Hospital, Athens, Greece | ²²⁸Riga East University Hospital, Riga, Latvia | ²²⁹Immunology and Allergy Division, Clinical Hospital, University of Chile, Santiago, Chile | ²³⁰Quality Use of Research Medicines Group, Woolcock Institute of Medical Research, Macquarie Park NSW, Australia | ²³¹Macquarie University, Macquarie Park, NSW, Australia | ²³²Faculty of Health, University of Plymouth, Plymouth, UK | ²³³Department of Primary Care and Population Health, University of Nicosia Medical School, Nicosia, Cyprus | ²³⁴Otolaryngology Department, General Hospital of Kalamata, Kalamata, Greece | ²³⁵Department of Medical Sciences, Occupational and Environmental Medicine, Uppsala University, Uppsala, Sweden | ²³⁶Clinical Management Group, Woolcock Institute of Medical Research, Sydney, Australia | ²³⁷Sydney Pharmacy School, Faculty of Medicine and Health, University of Sydney, Sydney, New South Wales, Australia | ²³⁸Departmental Unit of Allergy, Clinical Immunology & Pneumology, Istituto Ospedaliero Fondazione Poliambulanza, Brescia, Italy | ²³⁹Allergy Unit, Department of Internal Medicine, University Hospital AOU delle Marche, Ancona, Italy | ²⁴⁰Department of Clinical and Molecular Sciences, Marche Polytechnic University, Ancona, Italy | ²⁴¹Department of Otorhinolaryngology, Head and Neck Surgery, Ankara University Medical School, Ankara, Turkey | ²⁴²Department of Otorhinolaryngology Head and Neck Surgery, Salzburg Paracelsus Medical University, Salzburg, Austria | ²⁴³Department of Otolaryngology, Head and Neck Surgery, Weill Cornell Medical College, Cornell University, New York, NY, USA | ²⁴⁴Allergy and Clinical Immunology Department, University Hospital Center "Mother Teresa", Tirana, Albania | ²⁴⁵UMF-University of Medicine and Pharmacy 'Carol Davila', Pneumology Department, National Institute of Pneumology 'Marius Nasta', Bucharest, Romania | ²⁴⁶Department of Allergy and Internal Medicine, Medical University of Bialystok, Bialystok, Poland | ²⁴⁷Department of Otolaryngology-Head and Neck Surgery, Johns Hopkins University, Baltimore, Maryland, USA | ²⁴⁸Department of Pharmacotherapy and Pharmaceutical Care, Faculty of Pharmacy, Medical University of Warsaw, Warsaw, Poland | ²⁴⁹Otorhinolaryngology and Neck Surgery Unit, San Juan de Dios National Hospital, San Miguel, El Salvador | ²⁵⁰Ecole Polytechnique de Palaiseau, Palaiseau, France | ²⁵¹IRBA (Institut de Recherche Bio-Médicale Des Armées), Brétigny sur Orge, France | ²⁵²Université Paris Cité, Paris, France | ²⁵³Servicio de Alergia e Inmunología clínica, Hospital Universitario de Puebla, Puebla, México | ²⁵⁴Department of Otolaryngology-Head and Neck Surgery, Eye and Ear University Hospital, Beirut, Lebanon | ²⁵⁵Department of Otorhinolaryngology-Head and Neck Surgery, Dar Al Shifa Hospital, Salmiya, Kuwait | ²⁵⁶Center Allergy & Immunology and Geomedi Teaching University, Faculty of Medicine, Tbilisi, Georgia | ²⁵⁷Department of Otorhinolaryngology, Head and Neck Surgery, Interdisciplinary Graduate School of Medicine, University of Yamanashi, Yamanashi, Japan | ²⁵⁸Mohammed Bin Rashid University (MBRU), Dubai Health, Dubai, UAE | ²⁵⁹Asthma Reference Center—School of Medicine of Santa Casa de Misericórdia of Vitória, Espírito Santo, Brazil | ²⁶⁰ORL SOKLIC KOSAK, Ljubljana,

Slovenia | ²⁶¹Otolaryngology-HNS, University of Zurich, University Hospital of Zurich, Zurich, Switzerland | ²⁶²Head of Laboratory of Atmospheric Processes and Aerobiology, University of Latvia, Riga, Latvia | ²⁶³Clinic of Allergy and Immunology, University Clinical Center of Serbia, Belgrade, Serbia | ²⁶⁴Faculty of Medicine, University of Belgrade, Belgrade, Serbia | ²⁶⁵Department of Immunology and Allergy, University Hospital and Faculty of Medicine in Pilsen, Czech Republic | ²⁶⁶Charles University, Prague, Czech Republic | ²⁶⁷Department of Medicine Solna, Division of Immunology and Respiratory Medicine, Karolinska Institutet, Stockholm, Sweden | ²⁶⁸Department of Clinical Immunology and Transfusion Medicine, Karolinska University Hospital, Stockholm, Sweden | ²⁶⁹Center for Molecular Medicine, Karolinska University Hospital, Stockholm, Sweden | ²⁷⁰Department of Chest Diseases, Istanbul University-Cerrahpaşa, Cerrahpaşa Faculty of Medicine, Istanbul, Turkey | ²⁷¹Division of Allergy and Immunology, Department of Pediatrics, Siriraj Hospital, Mahidol University Faculty of Medicine, Bangkok, Thailand | ²⁷²Samitivej Allergy Institute, Bangkok, Thailand | ²⁷³Department of Otorhinolaryngology, Heinrich Heine University Düsseldorf, Medical Faculty and University Hospital Düsseldorf, Düsseldorf, Germany | ²⁷⁴Department of Medicine and Therapeutics, The Chinese University of Hong Kong, Hong Kong, China | ²⁷⁵ENT and Allergy, Düsseldorf, Germany | ²⁷⁶Shanghai Skin Disease Hospital, Tongji University School of Medicine, Shanghai, China

Acknowledgements

Open Access funding enabled and organized by Projekt DEAL.

Conflicts of Interest Management

Conflicts of interest (CoIs) of all participants were managed according to the Guidelines International Network (GIN) CoI declaration and management principles reviewed by the AWMF (*Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften e. V.*) [87]. A CoI was defined as an individual having "current material interests outside of ARIA that could influence or could be perceived as influencing his/her decisions, actions, or presentations."

Before participation, all guideline panel members and members of the evidence-synthesis team completed a standardised disclosure-of-interests (DoI) form provided by the AWMF. This form collected information about current and previous (past 3 years) interests in the healthcare sector, including direct financial interests and indirect (e.g., non-financial and intellectual) interests. Guideline panel members were required to report all interests in the DoI form, regardless of whether they would see such interests as having a thematic reference to the guideline or representing a conflict of interest.

Staff providing administrative support to ARIA-EAACI 2024–2025 judged every DoI form in the assessment on whether a disclosed interest was considered a CoI. The completed DoI forms are to be made available to all panel members during the project. At every meeting, guideline panellists were reminded of the DoI forms and prompted to make new disclosures. ARIA staff and members reviewed new disclosures, but previously documented disclosures or judgments are never removed or changed. The forms will receive a final update prior to guideline submission and will be included with the published guidelines.

The guideline panel was composed so that most of its members did not have a CoI. Absence of CoI is required for those chairing the panel discussions and gathering and summarising evidence for the EtDs. No individual was excluded from the guideline panel due to direct or indirect CoI. However, only those without CoI were allowed to make judgements or participate in the voting process for judgements or recommendations.

Conflicts of Interest

J. Bousquet reports personal fees from Cipla, Menarini, Mylan, Novartis, Purina, Sanofi-Aventis, Teva, Noucor, other from KYomed-Innov, other from Mask-air-SAS, outside the submitted work. M. Hyland reports

grants from GSK, from AstraZeneca, from null, outside the submitted work. A. Cruz reports grants, personal fees and non-financial support from AstraZeneca, personal fees from Chiesi, grants and personal fees from Eurofarma, personal fees from Farmoquímica, grants from EMS, grants and personal fees from GSK, grants and personal fees from Sanofi, outside the submitted work. J. Sastre reports grants and personal fees from Sanofi, personal fees from GSK, personal fees from Novartis, personal fees from AstraZeneca, personal fees from Mundipharma and personal fees from Faes Farma, outside the submitted work. D. Wallace reports as an author on the JTFPP practice parameters on Rhinitis, the most recent being 2020 and I have been an author on several previous ARIA and MASK-related articles. M. Wagenmann reports personal fees from Allergopharma, personal fees from ALK-Abello, grants and personal fees from AstraZeneca, personal fees from CSL Behring, grants and personal fees from GSK, personal fees from HAL Allergy, personal fees from Leti Pharma, personal fees from M.S.D., personal fees from Novartis, grants and personal fees from Regeneron, grants and personal fees from Sanofi, personal fees from Stallergenes, grants from Takeda, outside the submitted work. O. Pfaar reports grants and/or personal fees and/or travel support from AEDA, Alfried Krupp Krankenhaus, ALK-Abelló, Allergopharma, Almirall, Altamira Therapeutics, ASIT Biotech, AstraZeneca, Bencard Allergie GmbH/Allergy Therapeutics, Blueprint, Cliantha, Deutsche AllergieLigae.V., Deutsche Forschungsgemeinschaft, Dustri-Verlag, ECM Expro & Conference Management GmbH, Forum für Medizinische Fortbildung, Georg-Thieme-Verlag, GSK, HAL Allergy Holding B.V./HAL Allergie GmbH, Immunotek, Ingress Health, Institut für Disease Management (Essen, Germany), IQVIA Commercial, Japanese Society of Allergology, Königlich Dänisches Generalkonsulat, Laboratorios LETI/LETI Pharma, Lilly, Lofarma, Medizinische Hochschule Hannover, med update europe GmbH, Meinhardt Congress GmbH, Novartis, Paul-Ehrlich-Institut, Paul-Martini-Stiftung, PneumoLive, Pohl-Boskamp, Procter & Gamble, Red Maple Trials Inc., R.G. Aertzefortbildung, ROXALL Medizin, Sanofi-Aventis, Sanofi Genzyme, Springer Publisher, Stallergenes Greer, streamedup! GmbH, Technical University Dresden, Wiley Publishers, Wort & Bild Verlag, Verlag ME; outside the submitted work, Oliver Pfaar is Vice President of the European Academy of Allergy and Clinical Immunology (EAACI), a member of EAACI Excom as well as a member of the external board of directors of the German Society of Allergy and Clinical Immunology (DGAKI); coordinator, main- or co-author of different position papers and guidelines in rhinology, allergology and allergen immunotherapy; and he is Editor-in-Chief of Clinical Translational Allergy and Associate Editor of Allergy. T. Iinuma reports a grant from Sanofi, outside the submitted work. J. Davies reports grants from Australian National Health and Medical Research Council, grants from Australian Research Council, non-financial support from Sullivan Nicolaides Pathology Queensland, non-financial support from ThermoFisher Sweden, non-financial support from Swisens SA Switzerland, non-financial support from Kenelec Australia, non-financial support from Abacus Dx Australia, grants from Bayer Healthcare, grants from Australian Department of Health and Ageing, grants from Australian Medical Research Future Fund, outside the submitted work; in addition, Dr. Davies has a patent US PTO 14/311944 issued, a patent AU2008/316301 issued and a patent PCT/AU2024/051221 pending. I. Tsiligianni reports grants from Chiesi, GSK Hellas, Menarini, AstraZeneca Greece, outside the submitted work. B. Gradauskiene reports grants from AstraZeneca, personal fees from Berlin-Chemie Menarini, personal fees from Viartis, outside the submitted work. I. Ansotegui reports personal fees from Bayer, personal fees from Eurodrug, personal fees from Gebro, personal fees from Menarini, personal fees from M.S.D., personal fees from Roxall, personal fees from Sanofi, personal fees from Cipla, personal fees from Glenmark, personal fees from Opella, outside the submitted work. A. Fiocchi reports Danone Advisory board member, Abbott—Consultant, paid to institution, Aimmune—Advisory board member, Ferrero—Consultant, Novartis—Advisory board member, Sanofi—Advisory board member, Stallergenes—Advisory board member. M. Kupczyk reports personal fees from Adamed, personal fees from Abbvie, personal fees and non-financial support from AstraZeneca, personal fees and non-financial support from Berlin-Chemie, personal fees and

non-financial support from Chiesi, personal fees from EMMA, personal fees from GSK, personal fees from HAL Allergy, personal fees from Sunpharm, personal fees from HVD, personal fees from LEK-AM, personal fees from Polpharma, personal fees from Teva, personal fees from Sanofi, personal fees from Zentiva, personal fees from Stada, outside the submitted work. JC. Ivancevich reports personal fees from Laboratorios Casasco Argentina, outside the submitted work. A. Todo-Bom reports personal fees from GSK, grants and personal fees from LETI, personal fees from Mylan, grants from AbbVie, grants from Sanofi and grants from Roxal, outside the submitted work. L. Tuyet reports grants from AstraZeneca, grants from Boehringer Ingelheim, grants from Glaxo Smith Kline, grants from M.S.D., grants from Hyphens, grants from Tedis, grants from Novartis and grants from Orient Pharma, outside the submitted work. N. Papadopoulos reports grants from Nestle, Numil, Vianex, Vibrant, personal fees from Abbott, AstraZeneca, GSK, HAL, Menarini, Novartis, Danone Nutricia, OM Pharma, Regeneron, Sanofi, outside the submitted work. S. Toppila-Salmi reports grants and other support from Sanofi Pharma, grants and other support from GSK, other support from OrionPharma, other support from ALK-Abelló, other support from AstraZeneca, other support from Clario, outside the submitted work. R. Naclerio reports personal fees from Sanofi, personal fees from Lyra, outside the submitted work. M. Soyka reports other from Sanofi, other from M.S.D., other from GSK, other from AstraZeneca, outside the submitted work. N. Roche reports grants and personal fees from Boehringer Ingelheim, grants and personal fees from Novartis, grants and personal fees from GSK, personal fees from AstraZeneca, personal fees from Chiesi, grants and personal fees from Pfizer, personal fees from Sanofi, personal fees from Zambon, personal fees from M.S.D., personal fees from Austral, personal fees from Biosency, outside the submitted work. C. Suppli Ulrik reports personal fees from AZ, personal fees from GSK, grants and personal fees from BI, personal fees from TEVA, personal fees from Pfizer, personal fees from Orion, grants and personal fees from Sanofi, personal fees and non-financial support from Novartis, personal fees from Chiesi, personal fees from Covis Pharma, personal fees from Berlin-Chemie, outside the submitted work. D. Chu reports being a GRADE Working Group member, supports the development of allergy guidelines internationally and holds an E.J. Moran Campbell Career Award and a CIHR Inclusive Excellence Prize in Patient Engagement. Y. Okamoto reports personal fees from Torii Pharmaceutical Co. Ltd., personal fees from Tanabe Mitsubishi Pharmaceutical Co. Ltd., personal fees from Kirin Holdings Co. Ltd., personal fees from Shionogi Co. Ltd., personal fees from Stallergenes Greer and personal fees from Diichi-Sankyo, outside the submitted work. H. Olze received fees (lectures, advisory boards, research grants) from F. Hoffmann-La Roche Ltd., Sanofi-Aventis Deutschland GmbH, AstraZeneca GmbH, GlaxoSmithKline GmbH and Co. KG and Novartis, outside the present work. J. Mullol reports personal fees and other from Sanofi-Genzyme and Regeneron, grants, personal fees and other from Viartis/Meda Pharma, grants and personal fees from Noucor/Uriach Group, personal fees from Menarini, personal fees from UCB, personal fees and other from AstraZeneca, grants, personal fees and other from GSK, personal fees from M.S.D., personal fees and other from Lilly, personal fees and other from Glenmark, outside the submitted work. M. Torres reports personal fees from Leti Laboratories, personal fees from Aimmune Therapeutics, personal fees from Diater Laboratories, grants from European Commission, grants from SEAIC, grants from ISCHII, outside the submitted work. H. Kraxner reports Speaker's fee and congress support from Sanofi, Speaker's fee and congress support from Viartis, Speaker's fee and congress support from Berlin-Chemie, Speaker's fee and congress support from Ewopharma; Advisory Board membership: Sanofi. M. Makris reports personal fees from ELPEN, personal fees from MENARINI, personal fees from AstraZeneca, personal fees from Novartis, personal fees from CHIESI, personal fees from GSK, personal fees from ThermoFisher, personal fees from Vianex, outside the submitted work. D. Sakurai reports grants and personal fees from Torii, grants and personal fees from Tanabe Mitsubishi, personal fees from Shionogi, grants and personal fees from Taiho, grants and personal fees from Kyorin, personal fees from Meiji Seika Pharma, personal fees from Thermo Fisher Scientific Diagnostics, grants and personal fees from Tsumura, personal fees from Hisamitsu, personal

fees from Novartis, personal fees from Sanofi, personal fees from AstraZeneca, outside the submitted work. J. Correia-de-Sousa reports other from Boehringer Ingelheim, personal fees and other from GSK, grants, personal fees and other from AstraZeneca, personal fees from Bial, non-financial support from Mundipharma, personal fees from Sanofi, other from Novartis, personal fees from M.S.D., personal fees from Medinfar, outside the submitted work. M. Bourgoign-Heck reports non-financial support from ThermoFisher, personal fees from ALK, personal fees from Takeda, personal fees from Biocryst, personal fees from DBV, outside the submitted work. O. Palomares received research grants from MINECO, Ministerio de Ciencia e Innovación, CAM, Immunotek S.L., Novartis and AstraZeneca and fees for giving scientific lectures or participation in Advisory Boards from AstraZeneca, Pfizer, GlaxoSmithKline, Immunotek S.L., Novartis and Sanofi-Genzyme. M. Moniuszko reports having received in the past personal fees and other support from Berlin-Chemie/Menarini, personal fees and other support from AstraZeneca, personal fees and other support from GlaxoSmithKline, personal fees and other support from Novartis, personal fees and other support from Chiesi, personal fees and other support from Celon Pharma, personal fees and other support from Takeda, personal fees and other support from Polfarmex, personal fees and other support from CSL Behring, personal fees from Glenmark Pharmaceuticals, personal fees and other support from Sanofi, outside the submitted work. D. Larenas Linnemann reports personal fees from ALK, Armstrong, Astrazeneca national and global, Bayer, Chiesi, Grunenthal, Grin, GSK national and global, Viatriis, Menarini, M.S.D., Novartis, Pfizer, Sanofi, Siegfried, Carnot, Syneos Health, grants from Abbvie, Bayer, Lilly, Sanofi, Astrazeneca, Pfizer, Novartis, Pulmonair, GSK, Chiesi, Biopharma, outside the submitted work; and Editor-in-Chief of Immune System (Karger)—member of asthma committee ACAAI—subgroup chair of allergen immunotherapy Practice Parameter Update JTF AAAAI/ACAAI 2024—member of allergen immunotherapy committee AAAAI—chair of allergen immunotherapy committee CMICA—member of allergic asthma task force EAACI. M. Worm reports other from AbbVie Deutschland GmbH & Co. KG, other from Aimmune Therapeutics UK Limited, other from ALK-Abelló Arzneimittel GmbH, other from Allergopharma GmbH & Co KG, other from Almirall Hermal GmbH, other from Amgen GmbH, other from AstraZeneca GmbH, other from Bayer AG, other from Bencard Allergy GmbH, other from Bioprojet Pharma, other from Boehringer Ingelheim Pharma GmbH & Co. KG, other from Bristol Myers Squibb GmbH & Co. KGaA, other from Galderma Laboratorium GmbH, other from Glaxo Smith Kline GmbH & Co. KG, other from Infectopharm Arzneimittel und Consilium GmbH, other from LEO Pharma GmbH, other from Lilly Deutschland GmbH, other from Mylan Germany GmbH (A Viatriis Company), other from Novartis AG, other from Octapharm AG, other from Pfizer Pharma GmbH, other from Sanofi-Aventis Deutschland GmbH/Denzyme Europe B. B., outside the submitted work. T. Kosak Soklic reports personal fees from Meda AB, outside the submitted work. P. Devillier reports personal fees from ALK-Abello, personal fees and non-financial support from Stallergenes, personal fees and non-financial support from AstraZeneca, personal fees from Chiesi, personal fees from GlaxoSmithKline, personal fees from Menarini, personal fees from Viatriis, personal fees from Procter & Gamble Health, outside the submitted work. L. Cecchi reports personal fees from Menarini, personal fees from Chiesi, personal fees from AstraZeneca, personal fees from GSK, personal fees from Novartis, personal fees from ThermoFisher, personal fees from Sanofi, outside the submitted work. G. Roberts reports grants from the National Institute of Health Research, grants from the National Institutes of Health, grants from Action Medical Research, grants from EU IMI2, other support from ALK-Abello, non-financial support and other support from ThermoFisher, other support from AstraZeneca, other support from Viatriis, outside the submitted work. E. Compalati reports personal fees from Lofarma spa during the conduct of the study. D. Choudhury reports personal fees from Athaneum, others from ESPGHAN, others from the British Society of Allergy and Clinical Immunology (BSACI) - Standards of Care Committee, others from Respiratory Allergy, International Primary Care Respiratory Group, outside the submitted work. R. Buhl reports personal fees from AstraZeneca, Berlin-Chemie, Celltrion,

Chiesi, Cipla, Sanofi and Teva, as well as grants to Mainz University Hospital and personal fees from Boehringer Ingelheim, GlaxoSmithKline, Novartis and Roche, outside the submitted work. L. de las Vecillas reports fees for Lectures/Advisory Boards from AstraZeneca, GSK, LETI labs and Allergy Therapeutics, outside the submitted work. She has received a Research grant from the Spanish Society of Allergy and Clinical Immunology Foundation and from the ISCIII, outside the submitted work. L. Taborda-Barata reports personal fees from LETI, personal fees from Sanofi and personal fees from Diater, outside the submitted work. J. Bernstein reports personal fees from Pfizer, personal fees from Lilly and personal fees from Sanofi. S. Del Giacco reports grants and personal fees from AstraZeneca, grants and personal fees from GSK, grants and personal fees from Sanofi, personal fees from Chiesi, personal fees from Menarini, outside the submitted work. M. Giovannini reports personal fees from Sanofi, from Thermo Fisher Scientific, outside the submitted work. N. Miculinic reports personal fees from Salvus d.o.o. Pharma Industry, personal fees from Viatriis Ltd., outside the submitted work. J. da Silva reports grants, personal fees and others from Novartis, grants and personal fees from Takeda, outside the submitted work. G. Paoletti reports other from LoFarma, other from GSK and other from Astrazeneca, outside the submitted work. H. Schünemann reports and develops guidelines on Allergic Rhinitis in Asthma (ARIA), and his academic institution received research funding for it. The other authors declare no conflicts of interest, outside the submitted work.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

References

1. J. Bousquet, J. M. Anto, C. Bachert, et al., "Allergic Rhinitis," *Nature Reviews. Disease Primers* 6, no. 1 (2020): 95, <https://doi.org/10.1038/s41572-020-00227-0>.
2. S. Maio, S. Baldacci, L. Carrozzi, et al., "18-Yr Cumulative Incidence of Respiratory/Allergic Symptoms/Diseases and Risk Factors in the Pisa Epidemiological Study," *Respiratory Medicine* 158 (2019): 33–41.
3. M. Savouré, J. Bousquet, J. J. Jaakkola, M. S. Jaakkola, B. Jacquemin, and R. Nadif, "Worldwide Prevalence of Rhinitis in Adults: A Review of Definitions and Temporal Evolution," *Clinical and Translational Allergy* 12, no. 3 (2022): e12130.
4. T. Zuberbier, J. Lotvall, S. Simons, S. V. Subramanian, and M. K. Church, "Economic Burden of Inadequate Management of Allergic Diseases in the European Union: A GA(2) LEN Review," *Allergy* 69, no. 10 (2014): 1275–1279, <https://doi.org/10.1111/all.12470>.
5. J. Bousquet, P. Van Cauwenberge, N. Khaltaev, G. Aria Workshop, and O. World Health, "Allergic Rhinitis and Its Impact on Asthma," *Journal of Allergy and Clinical Immunology* 108, no. 5 Suppl (2001): S147–S334, <https://doi.org/10.1067/mai.2001.118891>.
6. J. L. Brozek, J. Bousquet, I. Agache, et al., "Allergic Rhinitis and Its Impact on Asthma (ARIA) Guidelines-2016 Revision," *Journal of Allergy and Clinical Immunology* 140, no. 4 (2017): 950–958, <https://doi.org/10.1016/j.jaci.2017.03.050>.
7. J. L. Brozek, J. Bousquet, C. E. Baena-Cagnani, et al., "Allergic Rhinitis and Its Impact on Asthma (ARIA) Guidelines: 2010 Revision," *Journal of Allergy and Clinical Immunology* 126, no. 3 (2010): 466–476, <https://doi.org/10.1016/j.jaci.2010.06.047>.
8. R. Brignardello-Petersen, A. Carrasco-Labra, and G. H. Guyatt, "How to Interpret and Use a Clinical Practice Guideline or Recommendation: Users' Guides to the Medical Literature," *Journal of the American Medical Association* 326, no. 15 (2021): 1516–1523, <https://doi.org/10.1001/jama.2021.15319>.
9. J. Bousquet, H. J. Schunemann, A. Togias, et al., "Next-Generation Allergic Rhinitis and Its Impact on Asthma (ARIA) Guidelines for

- Allergic Rhinitis Based on Grading of Recommendations Assessment, Development and Evaluation (GRADE) and Real-World Evidence," *Journal of Allergy and Clinical Immunology* 145, no. 1 (2020): 70–80.e3, <https://doi.org/10.1016/j.jaci.2019.06.049>.
10. J. Bousquet, H. J. Schunemann, B. Sousa-Pinto, et al., "Concepts for the Development of Person-Centered, Digitally Enabled, Artificial Intelligence-Assisted ARIA Care Pathways (ARIA 2024)," *Journal of Allergy and Clinical Immunology. In Practice* 12, no. 10 (2024): 2648–2668, <https://doi.org/10.1016/j.jaip.2024.06.040>.
11. P. Alonso-Coello, H. J. Schunemann, J. Moher, et al., "GRADE Evidence to Decision (EtD) Frameworks: A Systematic and Transparent Approach to Making Well Informed Healthcare Choices. 1: Introduction," *BMJ* 353 (2016): i2016, <https://doi.org/10.1136/bmj.i2016>.
12. P. Alonso-Coello, A. D. Oxman, J. Moher, et al., "GRADE Evidence to Decision (EtD) Frameworks: A Systematic and Transparent Approach to Making Well Informed Healthcare Choices. 2: Clinical Practice Guidelines," *BMJ* 353 (2016): i2089, <https://doi.org/10.1136/bmj.i2089>.
13. J. Bousquet, B. Sousa-Pinto, J. M. Anto, et al., "MASK-Air: An OECD (Organisation for Economic Co-Operation and Development) Best Practice for Public Health on Integrated Care for Chronic Diseases," *Journal of Allergy and Clinical Immunology. In Practice* 12, no. 8 (2024): 2010–2016, <https://doi.org/10.1016/j.jaip.2024.03.024>.
14. J. Bousquet, H. J. Schunemann, P. W. Hellings, et al., "MACVIA Clinical Decision Algorithm in Adolescents and Adults With Allergic Rhinitis," *Journal of Allergy and Clinical Immunology* 138, no. 2 (2016): 367–374.e2, <https://doi.org/10.1016/j.jaci.2016.03.025>.
15. J. Bousquet, J. M. Anto, B. Sousa-Pinto, et al., "Digitally-Enabled, Patient-Centred Care in Rhinitis and Asthma Multimorbidity: The ARIA-MASK-Air((R)) Approach," *Clinical and Translational Allergy* 13, no. 1 (2023): e12215, <https://doi.org/10.1002/ct2.12215>.
16. B. Sousa-Pinto, R. J. Vieira, M. Marques-Cruz, et al., "Artificial Intelligence-Supported Development of Health Guideline Questions," *Annals of Internal Medicine* 177 (2024): 1518–1529, <https://doi.org/10.7326/ANNALS-24-00363>.
17. P. Finnegan, M. Murphy, and C. O'Connor, "#Corticophobia: A Review on Online Misinformation Related to Topical Steroids," *Clinical and Experimental Dermatology* 48, no. 2 (2023): 112–115, <https://doi.org/10.1093/ced/llac019>.
18. B. Sousa-Pinto, A. Sa-Sousa, R. J. Vieira, et al., "Behavioural Patterns in Allergic Rhinitis Medication in Europe: A Study Using MASK-Air((R)) Real-World Data," *Allergy* 77, no. 9 (2022): 2699–2711, <https://doi.org/10.1111/all.15275>.
19. J. Bousquet, L. Klimek, P. Kuna, J. Mullol, and S. Toppila-Salmi, "The Debate: Regular Versus As-Needed Use of Intranasal Corticosteroids for a Patient-Centered Approach," *Journal of Allergy and Clinical Immunology. In Practice* 9, no. 3 (2021): 1374–1375, <https://doi.org/10.1016/j.jaip.2020.11.019>.
20. G. W. Canonica, F. Fumagalli, L. Guerra, et al., "Levocetirizine in Persistent Allergic Rhinitis: Continuous or On-Demand Use? A Pilot Study," *Current Medical Research and Opinion* 24, no. 10 (2008): 2829–2839, <https://doi.org/10.1185/03007990802395927>.
21. M. S. Dykewicz, H. B. Kaiser, R. A. Nathan, et al., "Fluticasone Propionate Aqueous Nasal Spray Improves Nasal Symptoms of Seasonal Allergic Rhinitis When Used as Needed (Prn)," *Annals of Allergy, Asthma & Immunology* 91, no. 1 (2003): 44–48, [https://doi.org/10.1016/S1081-1206\(10\)62057-1](https://doi.org/10.1016/S1081-1206(10)62057-1).
22. E. F. Juniper, G. H. Guyatt, P. M. O'Byrne, and M. Viveiros, "Aqueous Beclomethasone Dipropionate Nasal Spray: Regular Versus "as Required" Use in the Treatment of Seasonal Allergic Rhinitis," *Journal of Allergy and Clinical Immunology* 86, no. 3 Pt 1 (1990): 380–386, [https://doi.org/10.1016/s0091-6749\(05\)80101-0](https://doi.org/10.1016/s0091-6749(05)80101-0).
23. S. M. Kaszuba, F. M. Baroody, M. deTineo, L. Haney, C. Blair, and R. M. Naclerio, "Superiority of an Intranasal Corticosteroid Compared With an Oral Antihistamine in the As-Needed Treatment of Seasonal Allergic Rhinitis," *Archives of Internal Medicine* 161, no. 21 (2001): 2581–2587, <https://doi.org/10.1001/archinte.161.21.2581>.
24. P. Phinyo, C. Wongsas, M. Sompornrattanaphan, and T. Thongnarm, "As-Needed Versus Regular Intranasal Corticosteroid for Allergic Rhinitis: A Systematic Review and Meta-Analysis," *Asian Pacific Journal of Allergy and Immunology* 40, no. 3 (2022): 195–204, <https://doi.org/10.12932/AP-091121-1269>.
25. T. Thongnarm, C. Wongsas, P. Phinyo, P. Assanasen, P. Tantilipikorn, and M. Sompornrattanaphan, "As-Needed Versus Regular Use of Fluticasone Furoate Nasal Spray in Patients With Moderate to Severe, Persistent, Perennial Allergic Rhinitis: A Randomized Controlled Trial," *Journal of Allergy and Clinical Immunology. In Practice* 9, no. 3 (2021): 1365–1373.e6, <https://doi.org/10.1016/j.jaip.2020.09.057>.
26. J. B. Wartna, A. M. Bohnen, G. Elshout, et al., "Symptomatic Treatment of Pollen-Related Allergic Rhinoconjunctivitis in Children: Randomized Controlled Trial," *Allergy* 72, no. 4 (2017): 636–644, <https://doi.org/10.1111/all.13056>.
27. A. Anto, B. Sousa-Pinto, W. Czarlewski, et al., "Automatic Market Research of Mobile Health Apps for the Self-Management of Allergic Rhinitis," *Clinical and Experimental Allergy* 52, no. 10 (2022): 1195–1207, <https://doi.org/10.1111/cea.14135>.
28. J. Bousquet, A. Bedbrook, W. Czarlewski, et al., "Guidance to 2018 Good Practice: ARIA Digitally-Enabled, Integrated, Person-Centred Care for Rhinitis and Asthma," *Clinical and Translational Allergy* 9 (2019): 16, <https://doi.org/10.1186/s13601-019-0252-0>.
29. B. Sousa-Pinto, P. Eklund, O. Pfaar, et al., "Validity, Reliability, and Responsiveness of Daily Monitoring Visual Analog Scales in MASK-Air," *Clinical and Translational Allergy* 11, no. 7 (2021): e12062, <https://doi.org/10.1002/ct2.12062>.
30. J. Bousquet, B. Sousa-Pinto, J. M. Anto, et al., "Concurrent Validity, Cut-Offs and Ability to Change of Patient-Reported Outcome Measures for Rhinitis and Asthma in MASK-Air((R))," *Clinical and Translational Allergy* 14, no. 9 (2024): e12390, <https://doi.org/10.1002/ct2.12390>.
31. R. J. Vieira, B. Sousa-Pinto, A. Cardoso-Fernandes, et al., "Control of Allergic Rhinitis and Asthma Test: A Systematic Review of Measurement Properties and COSMIN Analysis," *Clinical and Translational Allergy* 12, no. 9 (2022): e12194, <https://doi.org/10.1002/ct2.12194>.
32. M. C. Reilly, A. Tanner, and E. O. Meltzer, "Work, Classroom and Activity Impairment Instruments: Validation Studies in Allergic Rhinitis," *Clinical Drug Investigation* 11 (1996): 278–288.
33. M. C. Reilly, A. S. Zbrozek, and E. M. Dukes, "The Validity and Reproducibility of a Work Productivity and Activity Impairment Instrument," *Pharmacoeconomics* 4, no. 5 (1993): 353–365.
34. B. Sousa-Pinto, G. Louis, J. Rodrigues, et al., "Impairment of EQ-5D-5L Domains According to Allergic Rhinitis and Asthma Control: A MASK-Air Real-World Study," *Journal of Allergy and Clinical Immunology. In Practice* 11 (2023): 3742–3751.e9, <https://doi.org/10.1016/j.jaip.2023.08.006>.
35. R. J. Vieira, L. Leemann, H. J. Schunemann, et al., "Impact of Uncontrolled Symptoms on the Health-Related Quality of Life (EQ-5D-5L) of Patients With Allergic Rhinitis: A MASK-Air Study," *Clinical and Experimental Allergy* 54, no. 8 (2024): 634–637, <https://doi.org/10.1111/cea.14516>.
36. M. Sofiev, Y. Palamarchuk, A. Bedard, et al., "A Demonstration Project of Global Alliance Against Chronic Respiratory Diseases: Prediction of Interactions Between Air Pollution and Allergen Exposure-The Mobile Airways Sentinel Network-Impact of Air POLLution on Asthma and Rhinitis Approach," *Chinese Medical Journal* 133, no. 13 (2020): 1561–1567, <https://doi.org/10.1097/CM9.0000000000000916>.

37. B. Sousa-Pinto, Y. Palamarchuk, L. Leemann, et al., "From MASK-Air and SILAM to CATALYSE (Climate Action to Advance HealthY Societies in Europe)," *Journal of Investigational Allergology & Clinical Immunology* 34, no. 1 (2024): 12–19, <https://doi.org/10.18176/jiaci.0923>.
38. J. Brozek, E. Borowiack, E. Sadowska, et al., "Patients' Values and Preferences for Health States in Allergic Rhinitis-An Artificial Intelligence Supported Systematic Review," *Allergy* 79, no. 7 (2024): 1812–1830, <https://doi.org/10.1111/all.16100>.
39. M. S. Dykewicz, D. V. Wallace, D. J. Amrol, et al., "Rhinitis 2020: A Practice Parameter Update," *Journal of Allergy and Clinical Immunology* 146, no. 4 (2020): 721–767, <https://doi.org/10.1016/j.jaci.2020.07.007>.
40. G. H. Guyatt, A. D. Oxman, R. Kunz, et al., "GRADE Guidelines: 2. Framing the Question and Deciding on Important Outcomes," *Journal of Clinical Epidemiology* 64, no. 4 (Apr 2011): 395–400, <https://doi.org/10.1016/j.jclinepi.2010.09.012>.
41. H. J. Schunemann, D. Armstrong, A. Degl'innocenti, et al., "A Randomized Multicenter Trial to Evaluate Simple Utility Elicitation Techniques in Patients With Gastroesophageal Reflux Disease," *Medical Care* 42, no. 11 (2004): 1132–1142, <https://doi.org/10.1097/00005650-200411000-00013>.
42. B. Sousa-Pinto, A. Valiulis, E. Melen, et al., "Asthma and Rhinitis Control in Adolescents and Young Adults: A Real-World MASK-Air Study," *Pediatric Allergy and Immunology* 35, no. 2 (2024): e14080, <https://doi.org/10.1111/pai.14080>.
43. L. Taborda-Barata, M. T. Ventura, H. Blain, et al., "MASK-Air(R) Real-World Data in Respiratory Allergy in Old-Age Adults," *Clinical and Translational Allergy* 13, no. 1 (2023): e12216, <https://doi.org/10.1002/ct2.12216>.
44. C. S. Rosario, C. A. Cardozo, H. J. C. Neto, and N. A. R. Filho, "Do Gender and Puberty Influence Allergic Diseases?," *Allergologia et Immunopathologia* 49, no. 2 (2021): 122–125, <https://doi.org/10.15586/aei.v49i2.49>.
45. T. Keller, C. Hohmann, M. Standl, et al., "The Sex-Shift in Single Disease and Multimorbid Asthma and Rhinitis During Puberty—A Study by MeDALL," *Allergy* 73, no. 3 (2018): 602–614, <https://doi.org/10.1111/all.13312>.
46. K. Goldhahn, A. Bockelbrink, M. Nocon, et al., "Sex-Specific Differences in Allergic Sensitization to House Dust Mites: A Meta-Analysis," *Annals of Allergy, Asthma & Immunology* 102, no. 6 (2009): 487–494.
47. K. K. Gupta and S. Anari, "Medical Management of Rhinitis in Pregnancy," *Auris, Nasus, Larynx* 49, no. 6 (2022): 905–911, <https://doi.org/10.1016/j.anl.2022.01.014>.
48. J. Bousquet, E. Melen, T. Haahtela, et al., "Rhinitis Associated With Asthma Is Distinct From Rhinitis Alone: The ARIA-MeDALL Hypothesis," *Allergy* 78, no. 5 (2023): 1169–1203, <https://doi.org/10.1111/all.15679>.
49. J. M. Anto, J. Bousquet, M. Akdis, et al., "Mechanisms of the Development of Allergy (MeDALL): Introducing Novel Concepts in Allergy Phenotypes," *Journal of Allergy and Clinical Immunology* 139, no. 2 (2017): 388–399, <https://doi.org/10.1016/j.jaci.2016.12.940>.
50. J. Bousquet, B. Sousa-Pinto, F. S. Regateiro, et al., "MASK-Air Direct Patient Data Support the ARIA-MeDALL Hypothesis on Allergic Phenotypes," *Allergy* 78, no. 10 (2023): 2790–2794, <https://doi.org/10.1111/all.15842>.
51. J. Bousquet, P. Devillier, J. M. Anto, et al., "Daily Allergic Multimorbidity in Rhinitis Using Mobile Technology: A Novel Concept of the MASK Study," *Allergy* 73, no. 8 (2018): 1622–1631, <https://doi.org/10.1111/all.13448>.
52. F. Cibella, G. Ferrante, G. Cuttitta, et al., "The Burden of Rhinitis and Rhinoconjunctivitis in Adolescents," *Allergy, Asthma & Immunology Research* 7, no. 1 (2015): 44–50, <https://doi.org/10.4168/aaair.2015.7.1.44>.
53. E. Burte, J. Bousquet, V. Siroux, J. Just, B. Jacquemin, and R. Nadif, "The Sensitization Pattern Differs According to Rhinitis and Asthma Multimorbidity in Adults: The EGEEA Study," *Clinical and Experimental Allergy* 47, no. 4 (2017): 520–529, <https://doi.org/10.1111/cea.12897>.
54. V. Siroux, A. Boudier, R. Nadif, C. Lupinek, R. Valenta, and J. Bousquet, "Association Between Asthma, Rhinitis, and Conjunctivitis Multimorbidities With Molecular IgE Sensitization in Adults," *Allergy* 74, no. 4 (2019): 824–827, <https://doi.org/10.1111/all.13676>.
55. R. Amaral, J. Bousquet, A. M. Pereira, et al., "Disentangling the Heterogeneity of Allergic Respiratory Diseases by Latent Class Analysis Reveals Novel Phenotypes," *Allergy* 74, no. 4 (2019): 698–708, <https://doi.org/10.1111/all.13670>.
56. F. Raciborski, J. Bousquet, J. Bousquet, et al., "Dissociating Polysensitization and Multimorbidity in Children and Adults From a Polish General Population Cohort," *Clinical and Translational Allergy* 9 (2019): 4, <https://doi.org/10.1186/s13601-019-0246-y>.
57. I. Neumann, J. M. Anto, J. Bousquet, and H. J. Schunemann, "The Impact of Climate Change on Health Needs Structured Evidence Assessment and an Evidence to Action Framework to Make Decisions: A Proposal to Adopt the GRADE Approach," *Journal of Clinical Epidemiology* 157 (2023): 146–153, <https://doi.org/10.1016/j.jclinepi.2023.01.006>.
58. D. K. Chu, R. Brignardello-Petersen, G. H. Guyatt, C. Ricci, and J. Genueneit, "Method's Corner: Allergist's Guide to Network Meta-Analysis," *Pediatric Allergy and Immunology* 33, no. 1 (2022): e13609, <https://doi.org/10.1111/pai.13609>.
59. B. Sousa-Pinto, R. J. Vieira, A. Bognanni, et al., "Efficacy and Safety of Intranasal Medications for Allergic Rhinitis: Network Meta-Analysis," *Allergy* 80, no. 1 (2025): 94–105.
60. B. Sousa-Pinto, R. J. Vieira, J. Brozek, et al., "Intranasal Antihistamines and Corticosteroids in Allergic Rhinitis: A Systematic Review and Meta-Analysis," *Journal of Allergy and Clinical Immunology* 154, no. 2 (2024): 340–354, <https://doi.org/10.1016/j.jaci.2024.04.016>.
61. M. I. Torres, S. Gil-Mata, A. Bognanni, et al., "Intranasal Versus Oral Treatments for Allergic Rhinitis: A Systematic Review With Meta-Analysis," *Journal of Allergy and Clinical Immunology. In Practice* 12 (2024): 3404–3418, <https://doi.org/10.1016/j.jaip.2024.09.001>.
62. B. Sousa-Pinto, R. J. Vieira, J. Brozek, et al., "Intranasal Antihistamines and Corticosteroids in the Treatment of Allergic Rhinitis: A Systematic Review and Meta-Analysis Protocol," *BMJ Open* 13, no. 11 (2023): e076614, <https://doi.org/10.1136/bmjopen-2023-076614>.
63. R. J. Vieira, M. I. Torres, A. Bognanni, et al., "Protocol for the Systematic Reviews on the Desirable and Undesirable Effects of Pharmacological Treatments of Allergic Rhinitis Informing the ARIA 2024 Guidelines," *Allergol Select* 8 (2024): 270–277, <https://doi.org/10.5414/ALX02515E>.
64. J. B. Kostis and J. M. Dobrzynski, "Limitations of Randomized Clinical Trials," *American Journal of Cardiology* 129 (2020): 109–115, <https://doi.org/10.1016/j.amjcard.2020.05.011>.
65. R. Tsang, L. Colley, and L. D. Lynd, "Inadequate Statistical Power to Detect Clinically Significant Differences in Adverse Event Rates in Randomized Controlled Trials," *Journal of Clinical Epidemiology* 62, no. 6 (2009): 609–616, <https://doi.org/10.1016/j.jclinepi.2008.08.005>.
66. U. M. Centre, "About VigiBase," <https://who-umc.org/vigibase/>.
67. K. J. Rothman, S. Lanes, and S. T. Sacks, "The Reporting Odds Ratio and Its Advantages Over the Proportional Reporting Ratio," *Pharmacoepidemiology and Drug Safety* 13, no. 8 (2004): 519–523, <https://doi.org/10.1002/pds.1001>.
68. R. J. Vieira, B. Sousa-Pinto, A. Bognanni, et al., "Embedding Patients' Values and Preferences in Guideline Development for Allergic Diseases: The Case Study of Allergic Rhinitis and Its Impact on Asthma 2024," *Clinical and Translational Allergy* 14, no. 6 (2024): e12377, <https://doi.org/10.1002/ct2.12377>.

69. R. J. Vieira, L. Leemann, A. Briggs, et al., "Poor Rhinitis and Asthma Control Is Associated With Decreased Health-Related Quality of Life and Utilities: A MASK-Air Study," *Journal of Allergy and Clinical Immunology. In Practice* 12, no. 6 (Jun 2024): 1530–1538 e6, <https://doi.org/10.1016/j.jaip.2024.03.036>.
70. S. Gil-Mata, T. Teixeira, A. Bedbrook, et al., "Perceptions of the Impact of Individual Allergic Rhinitis Symptoms: A Survey of ARIA Clinical Experts," *World Allergy Organization Journal* 17, no. 12 (2024): 100999, <https://doi.org/10.1016/j.waojou.2024.100999>.
71. R. J. Vieira, L. F. Azevedo, A. M. Pereira, et al., "Impact of Allergic Rhinitis Control on Work Productivity and Costs: A Real-World Data MASK-Air Study," *Journal of Allergy and Clinical Immunology. In Practice* 12 (2024): 3107–3115.e13, <https://doi.org/10.1016/j.jaip.2024.07.026>.
72. Health CaFDaTi, *CDR Pharmacoeconomic Review Report for Dymista* (2016), https://www.cda-amc.ca/sites/default/files/cdr/pharmaco-economic/SR0408_Dymista_PE_Report.pdf.
73. World Health Organization, *The Selection and Use of Essential Medicines: Report of the WHO Expert Committee on Selection and Use of Essential Medicines, 2023 (Including the 23rd WHO Model List of Essential Medicines and the 9th WHO Model List of Essential Medicines for Children)* (World Health Organization, 2024).
74. R. Shikhar and A. M. Rentz, "Satisfaction With Medication: An Overview of Conceptual, Methodologic, and Regulatory Issues," *Value in Health* 7, no. 2 (2004): 204–215, <https://doi.org/10.1111/j.1524-4733.2004.72252.x>.
75. B. Sousa-Pinto, H. J. Schunemann, A. Sa-Sousa, et al., "Comparison of Rhinitis Treatments Using MASK-Air(R) Data and Considering the Minimal Important Difference," *Allergy* 77, no. 10 (2022): 3002–3014, <https://doi.org/10.1111/all.15371>.
76. E. Menditto, E. Costa, L. Midao, et al., "Adherence to Treatment in Allergic Rhinitis Using Mobile Technology the MASK Study," *Clinical and Experimental Allergy* 49, no. 4 (2019): 442–460, <https://doi.org/10.1111/cea.13333>.
77. B. Sousa-Pinto, E. M. Costa, R. J. Vieira, et al., "Adherence to Treatment in Allergic Rhinitis During the Pollen Season in Europe: A MASK-Air Study," *Clinical and Experimental Allergy* 55 (2025): 226–238, <https://doi.org/10.1111/cea.70004>.
78. European Medicines Agency, *Guideline on the Clinical Development of Medicinal Products for the Treatment of Allergic Rhinoconjunctivitis* (2004).
79. S. Whitmee, A. Haines, C. Beyrer, et al., "Safeguarding Human Health in the Anthropocene Epoch: Report of the Rockefeller Foundation–Lancet Commission on Planetary Health," *Lancet* 386, no. 10007 (2015): 1973–2028.
80. A. Herrmann, B. Lenzer, B. S. Müller, et al., "Integrating Planetary Health Into Clinical Guidelines to Sustainably Transform Health Care," *Lancet Planetary Health* 6, no. 3 (2022): e184–e185.
81. J. Bousquet, J. M. Anto, T. Haahtela, et al., "Digital Transformation of Health and Care to Sustain Planetary Health: The MASK Proof-of-Concept for Airway Diseases-POLLAR Symposium Under the Auspices of Finland's Presidency of the EU, 2019 and MACVIA-France, Global Alliance Against Chronic Respiratory Diseases (GARD, WH0) Demonstration Project, Reference Site Collaborative Network of the European Innovation Partnership on Active and Healthy Ageing," *Clinical and Translational Allergy* 10 (2020): 24, <https://doi.org/10.1186/s13601-020-00321-2>.
82. J. I. Halonen, M. Erhola, E. Furman, et al., "The Helsinki Declaration 2020: Europe That Protects," *Lancet Planetary Health* 4, no. 11 (2020): e503–e505, [https://doi.org/10.1016/S2542-5196\(20\)30242-4](https://doi.org/10.1016/S2542-5196(20)30242-4).
83. J. I. Halonen, M. Erhola, E. Furman, et al., "A Call for Urgent Action to Safeguard Our Planet and Our Health in Line With the Helsinki Declaration," *Environmental Research* 193 (2021): 110600, <https://doi.org/10.1016/j.envres.2020.110600>.
84. J. Andrews, G. Guyatt, A. D. Oxman, et al., "GRADE Guidelines: 14. Going From Evidence to Recommendations: The Significance and Presentation of Recommendations," *Journal of Clinical Epidemiology* 66, no. 7 (2013): 719–725, <https://doi.org/10.1016/j.jclinepi.2012.03.013>.
85. J. C. Andrews, H. J. Schunemann, A. D. Oxman, et al., "GRADE Guidelines: 15. Going From Evidence to Recommendation—Determinants of a Recommendation's Direction and Strength," *Journal of Clinical Epidemiology* 66, no. 7 (2013): 726–735, <https://doi.org/10.1016/j.jclinepi.2013.02.003>.
86. A. L. Courbis, R. B. Murray, S. Arnavielhe, et al., "Electronic Clinical Decision Support System for Allergic Rhinitis Management: MASK e-CDSS," *Clinical and Experimental Allergy* 48, no. 12 (2018): 1640–1653, <https://doi.org/10.1111/cea.13230>.
87. H. J. Schunemann, L. A. Al-Ansary, F. Forland, et al., "Guidelines International Network: Principles for Disclosure of Interests and Management of Conflicts in Guidelines," *Annals of Internal Medicine* 163, no. 7 (2015): 548–553.

Supporting Information

Additional supporting information can be found online in the Supporting Information section. **Appendix S1:** all70100-sup-0001-AppendixS1.docx.